

# Deutsches Patent- und Markenamt

München, den 1. August 2003


Telefon: (0 89) 21 95 - 4290

Aktenzeichen: 101 58 523.3-41

Deutsches Patent- und Markenamt · 80297 München

Anmelder: F. Hoffmann-La Roche AG

Patentanwälte  
Vossius & Partner  
Abhofach im DPMA  
München



Ihr Zeichen: F 2948 DE

Bitte Aktenzeichen und Anmelder bei  
allen Eingaben und Zahlungen angeben

Zutreffendes ist angekreuzt ☒ und/oder ausgefüllt

Prüfungsantrag, Einzahlungstag am 29. November 2001

Eingabe vom

eingegangen am

Die Prüfung der oben genannten Patentanmeldung hat zu dem nachstehenden Ergebnis geführt.

Zur Äußerung wird eine Frist von

**sechs Monat(en)**

gewährt, die mit der Zustellung beginnt.

Für Unterlagen, die der Äußerung gegebenenfalls beigelegt werden (z.B. Beschreibung, Beschreibungsteile, Patentansprüche, Zeichnungen), sind je zwei Ausfertigungen auf gesonderten Blättern erforderlich. Die Äußerung selbst wird nur in einfacher Ausfertigung benötigt.

Werden die Beschreibung, die Patentansprüche oder die Zeichnungen im Laufe des Verfahrens geändert, so hat der Anmelder, sofern die Änderungen nicht vom Deutschen Patent- und Markenamt vorgeschlagen sind, im Einzelnen anzugeben, an welcher Stelle die in den neuen Unterlagen beschriebenen Erfindungsmerkmale in den ursprünglichen Unterlagen offenbart sind.

☒ In diesem Bescheid sind folgende Entgegenhaltungen erstmalig genannt (Bei deren Nummerierung gilt diese auch für das weitere Verfahren):

(1) WO 00/63364 A2, s. bes. S. 20-21

## Hinweis auf die Möglichkeit der Gebrauchsmusterabzweigung

Der Anmelder einer mit Wirkung für die Bundesrepublik Deutschland eingereichten Patentanmeldung kann eine Gebrauchsmusteranmeldung, die den gleichen Gegenstand betrifft, einreichen und gleichzeitig den Anmeldetag der früheren Patentanmeldung in Anspruch nehmen. Diese Abzweigung (§ 5 Gebrauchsmustergesetz) ist bis zum Ablauf von 2 Monaten nach dem Ende des Monats möglich, in dem die Patentanmeldung durch rechtskräftige Zurückweisung, freiwillige Rücknahme oder Rücknahmefiktion erledigt, ein Einspruchsverfahren abgeschlossen oder - im Falle der Erteilung des Patents - die Frist für die Beschwerde gegen den Erteilungsbeschluss fruchtlos verstrichen ist. Ausführliche Informationen über die Erfordernisse einer Gebrauchsmusteranmeldung, einschließlich der Abzweigung, enthält das Merkblatt für Gebrauchsmusteranmelder (G 6181), welches kostenlos beim Deutschen Patent- und Markenamt und den Patentinformationszentren erhältlich ist.

Annahmestelle und  
Nachbriefkasten  
nur  
Zweibrückenstraße 12

Hauptgebäude  
Zweibrückenstraße 12  
Zweibrückenstraße 5-7 (Brallierhof)  
Markenabteilungen:  
Cincinnatistraße 64  
81534 München

Hausadresse (für Fracht)  
Deutsches Patent- und Markenamt  
Zweibrückenstraße 12  
80333 München

Telefon (089) 2195-0  
Telefax (089) 2195-2231  
Internet: <http://www.dpma.de>

Bank:  
88k München  
Kto.Nr.: 700 010 54  
BLZ: 700 000 00

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/11, 15/63, 15/70, 15/86, C07K 14/01, 14/08, A61K 48/00, 38/16, A61P 35/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/63364</b> <b>(43) International Publication Date:</b> 26 October 2000 (26.10.00)
<b>(21) International Application Number:</b> PCT/US00/10555 <b>(22) International Filing Date:</b> 19 April 2000 (19.04.00) <b>(30) Priority Data:</b> 60/130,377 21 April 1999 (21.04.99) US <b>(71) Applicant (for all designated States except US):</b> AMERICAN HOME PRODUCTS CORPORATION [US/US]; Five Giralda Farms, Madison, NJ 07940 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> PACHUK, Catherine [US/US]; 3212 New Park Circle, Lansdale, PA 19446 (US). SATISHCHANDRAN, C. [US/US]; 605 Shepard Drive, Lansdale, PA 19446 (US). <b>(74) Agents:</b> BAK, Mary, E. et al.; Howson and Howson, Spring House Corporate Center, P.O. Box 457, Spring House, PA 19477 (US).	<b>(81) Designated States:</b> AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>	
<b>(54) Title:</b> METHODS AND COMPOSITIONS FOR INHIBITING THE FUNCTION OF POLYNUCLEOTIDE SEQUENCES  <b>(57) Abstract</b>  A therapeutic composition for inhibiting the function of a target polynucleotide sequence in a mammalian cell includes an agent that provides to a mammalian cell an at least partially double-stranded RNA molecule comprising a polynucleotide sequence of at least about 200 nucleotides in length, said polynucleotide sequence being substantially homologous to a target polynucleotide sequence. This RNA molecule desirably does not produce a functional protein. The agents useful in the composition can be RNA molecules made by enzymatic synthetic methods or chemical synthetic methods <i>in vitro</i> ; or made in recombinant cultures of microorganisms and isolated therefrom, or alternatively, can be capable of generating the desired RNA molecule <i>in vivo</i> after delivery to the mammalian cell. In methods of treatment of prophylaxis of virus infections, other pathogenic infections or certain cancers, these compositions are administered in amounts effective to reduce or inhibit the function of the target polynucleotide sequence, which can be of pathogenic origin or produced in response to a tumor or other cancer, among other sources.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## METHODS AND COMPOSITIONS FOR INHIBITING THE FUNCTION OF POLYNUCLEOTIDE SEQUENCES

### Field of the Invention

The present invention relates to polynucleotide compositions which have an  
5 inhibitory or other regulatory effect upon the function of certain target polynucleotide  
sequences present in a mammalian cell, and for methods of using the compositions in  
therapeutic, prophylactic, diagnostic and research methods.

### Background of the Invention

Polynucleotide compositions have been described for pharmaceutical uses,  
10 primarily for treatment or prophylaxis of disease in mammals, as well as in research in  
such fields. Specifically a great deal of activity presently surrounds the use of  
polynucleotide compositions in the treatment of pathogenic extracellular and  
intracellular infections, such as viral, bacterial, fungal infections, and the like. As one  
example, DNA vaccines are described to deliver to a mammalian cell *in vivo* an agent  
15 which combats a pathogen by harnessing the mammalian immune system. Thus, such  
vaccines are designed to express, for example, a viral protein or polypeptide, and elicit  
a humoral or cellular immune response upon challenge by the infective agent. Gene  
therapy vectors, on the other hand, are polynucleotide compositions generally  
designed to deliver to a mammalian cell a protein which is either not expressed,  
20 expressed improperly or underexpressed in a mammal. Such vectors frequently must  
address species specific immune responses to the those polynucleotide sequences that  
are recognized as antigenic or which evoke an unwanted cellular immune response.

Still other therapeutic uses of polynucleotide compositions are for the delivery  
of missing or underexpressed proteins to a diseased mammalian patient. Furthermore,  
25 polynucleotides are useful themselves as *in vivo* reagents, in diagnostic/imaging  
protocols, as reagents in gene therapy, in antisense protocols and in vaccine  
applications or otherwise as pharmaceuticals used to treat or prevent a variety of  
ailments such as genetic defects, infectious diseases, cancer, and autoimmune diseases.

Polynucleotides are also useful as *in vitro* reagents in assays such as biological research assays, medical, diagnostic and screening assays and contamination detection assays.

5 A host of problems well-known to the art has prevented the numerous polynucleotide compositions from becoming widely accepted as useful pharmaceuticals. Thus, there are few such DNA vaccines or therapeutics which have yet been accepted by the medical community for the treatment of disease in mammals.

Phenomena have been observed in plants and nematodes that are mediated by polynucleotide compositions, and are referred to as post-transcriptional gene silencing and transcriptional silencing. This phenomenon demonstrates that transfection or  
10 infection of a plant, nematode or *Drosophila* with a virus, viroid, plasmid or RNA expressing a polynucleotide sequence having some homology to a regulatory element, such as a promoter or a native gene or a portion thereof already expressed in that cell, can result in the permanent inhibition of expression of both the endogenous regulatory element or gene and the exogenous sequence. This silencing effect was shown to be  
15 gene specific. See, for example, L. Timmons and A. Fire, Nature, 395:354 (Oct. 29, 1998); A. Fire *et al*, Nature, 391:806-810 (Feb. 19, 1998); and R. Jorgensen *et al*, Science, 279:1486-1487 (March 6, 1998)]. A DNA plasmid encoding a full-length pro-alpha 1 collagen gene was transiently transfected into a rodent fibroblast tissue cell line and a "silencing" effect on the native collagen gene and the transiently  
20 expressed gene observed [Bahramian and Zarbl, Mol. Cell. Biol., 19(1):274-283 (Jan. 1999)].

See, also, International Patent Application No. WO98/05770, published February 12, 1998, which relates to gene inhibition by use of an antisense RNA with  
25 secondary structures, and/or in combination with double stranded RNase. International Patent Application No. WO99/53050, published October 21, 1999, also relates to reducing phenotypic expression of a nucleic acid, particularly in plant cells, by introducing chimeric genes encoding sense and anti-sense RNA molecules.

There exists a need in the art for polynucleotide compositions and methods of  
30 using same to inhibit the function of polynucleotide sequences which are disease-

causing in mammals, such as polynucleotide sequences essential for the replication of viruses and other intracellular pathogens in mammalian cells, or sequences of extracellular mammalian pathogens, or sequences of tumor antigens which mediate the spread of cancer in a mammal, and the like, without adversely affecting essential gene sequences in the mammal.

### Summary of the Invention

In one aspect, the invention provides a composition for inhibiting the function of a target polynucleotide sequence in a mammalian cell. The composition comprises an agent that provides to a mammalian cell an at least partially double-stranded RNA molecule comprising a polynucleotide sequence of at least about 200 nucleotides in length. The polynucleotide sequence is substantially homologous to the target polynucleotide sequence, which can be a polynucleotide sequence, e.g., of a virus or other intracellular pathogen, a polynucleotide sequence of a cancer antigen or of an essential tumorigenic regulatory sequence, a polynucleotide sequence of an extracellular pathogen present in a mammal, or any other polynucleotide sequence which is desired to be "turned off" in a cell. This RNA molecule preferably does not produce a functional protein. This RNA molecule is preferably substantially non-homologous to naturally-occurring, essential mammalian polynucleotide sequences. In one embodiment, the agent of this composition is an RNA molecule made by enzymatic synthetic methods or chemical synthetic methods *in vitro*. In another embodiment, the RNA molecule may be generated in a recombinant culture, e.g., bacterial cells, isolated therefrom, and used in the methods discussed below. In another embodiment the agent of this composition generates the RNA molecule *in vivo* after delivery to the mammalian cell.

In another aspect, the invention provides a pharmaceutical composition comprising one or more of the compositions described immediately above and specifically hereinbelow, and an optional second agent that facilitates polynucleotide uptake in a cell, in a pharmaceutically acceptable carrier. Such compositions are useful for treating intracellular pathogenic infections, such as viruses. Other such

compositions are useful for treating certain cancers. Other such compositions are useful for treating certain extracellular pathogens. Still other such compositions are useful for treating any disease or disorder wherein inhibiting the function of a polynucleotide sequence in a mammal is desirable for therapy or vaccine use.

5           In still another aspect, the invention provides a method for treating a viral infection in a mammal by administering to the mammal one or more of the above-described compositions wherein the target polynucleotide is a virus polynucleotide sequence necessary for replication and/or pathogenesis of the virus in an infected mammalian cell, along with an optional second agent that facilitates polynucleotide uptake in a cell, in a pharmaceutically acceptable carrier. This composition is  
10 administered in an amount effective to reduce or inhibit the function of the viral sequence in the cells of the mammal.

          In yet a further aspect, the invention provides a method for preventing a viral infection in a mammal by administering to the mammal one or more of the above-  
15 described compositions wherein the target polynucleotide is a virus polynucleotide sequence necessary for replication and/or pathogenesis of the virus in an infected mammalian cell, with an optional second agent that facilitates polynucleotide uptake in a cell, in a pharmaceutically acceptable carrier. This composition is administered in an amount effective to reduce or inhibit the function of the viral sequence upon  
20 subsequent introduction of the virus into the mammalian cells.

          In still another aspect, the invention provides a method for treatment or prophylaxis of a virally induced cancer in a mammal by administering to the mammal one or more of the above described compositions in which the target polynucleotide is a sequence encoding a tumor antigen or functional fragment thereof or a regulatory  
25 sequence, which sequence function is required for the maintenance of the tumor in the mammal. The compositions can contain an optional second agent that facilitates polynucleotide uptake in a cell, and a pharmaceutically acceptable carrier. The composition is administered in an amount effective to reduce or inhibit the function of the tumor-maintaining sequence in the mammal.

In another aspect, the invention provides a method for the treatment or prophylaxis of infection of a mammal by an intracellular pathogen. The mammal is administered one or more of the compositions herein described wherein the target polynucleotide is a polynucleotide sequence of the intracellular pathogen necessary for replication and/or pathogenesis of the pathogen in an infected mammalian cell. The composition is administered with an optional second agent that facilitates polynucleotide uptake in a cell, in a pharmaceutically acceptable carrier, in an amount effective to reduce or inhibit the function of the sequence in the mammal.

In another aspect, the invention provides a method for the treatment or prophylaxis of infection of a mammal by an extracellular mammalian pathogen. The mammal is administered one or more of the compositions herein described wherein the target polynucleotide is a polynucleotide sequence of the extracellular pathogen necessary for replication and/or pathogenesis of the pathogen in an infected mammal. The composition is administered in a pharmaceutically acceptable carrier, in an amount effective to reduce or inhibit the function of the sequence in the mammal. It may be administered with an optional second agent that facilitates polynucleotide uptake by the pathogenic cell.

In still another aspect, the invention provides a method of treatment or prophylaxis of cancer in a mammal. The mammal is administered one or more of the above-described compositions, wherein the target polynucleotide is a polynucleotide sequence of an abnormal cancer-causing gene or non-expressed regulatory sequence in a mammal, which also possesses a normal copy of the gene or regulatory sequence. According to this aspect, the differences between the abnormal sequence and the normal sequence are differences in polynucleotides. The composition is administered with an optional second agent that facilitates polynucleotide uptake in a cell, in a pharmaceutically acceptable carrier, and in an amount effective to reduce or inhibit the function of the abnormal sequence in the mammal.

In yet a further aspect, the invention involves a method for treating a disease or disorder in a mammal comprising administering to the mammal having a disease or disorder characterized by expression of polynucleotide product not found in a healthy



mammal, one or more of the compositions as above described, in which the target polynucleotide sequence is the polynucleotide sequence which expresses that polynucleotide product or a non-expressed regulatory sequence essential to the expression of that product. The composition is administered with or without a second  
5 agent that facilitates polynucleotide uptake in a cell, and in a pharmaceutically acceptable carrier, in an amount effective to reduce or inhibit the function of the target polynucleotide product or regulatory sequence in the cells of the mammal.

Still another aspect of the present invention provides such compositions for use in research methods, such as a reagent for reducing or inhibiting undesired gene  
10 expression in mammalian cells or tissue *in vitro* for use in diagnostic or other research assays, or *ex vivo* for return to the mammal for therapy or other medicinal uses.

Other aspects of the invention are described further in the following detailed description of the preferred embodiments thereof.

#### **Brief Description of the Figures**

15 Fig. 1A is an illustration of a PCR product generated using the bacteriophage T7 RNA polymerase promoter- forward gag primer (T7F) and reverse gag (R) primer. Transcription from this PCR template, using T7 RNA polymerase generates an RNA sequence gag sense strand.

Fig. 1B is an illustration of a PCR product generated using a forward gag  
20 primer (F) and T7 promoter reverse gag (T7R) primer. Transcription of this template using a T7 RNA polymerase generates an RNA sequence gag antisense strand. Use of both the template of Fig. 1A and the template of Fig. 1B yields double stranded gag RNA sequence.

#### **Detailed Description of the Invention**

25 The present invention provides novel polynucleotide compositions and methods for therapy, prophylaxis, research and diagnostics in diseases and disorders which afflict mammalian species, in which the goal is to reduce or inhibit the function of a selected target polynucleotide sequence. These compositions and methods have

utility both *in vitro* and *in vivo*. These compositions and methods further enable the harnessing of the molecular mechanisms of the cell to accomplish therapeutic goals without requiring any stimulation of the immune system of the mammal involved.

As used herein, the phrases "target" or "target polynucleotide sequence" refer to any sequence present in a mammalian cell or in a mammalian organism, whether a naturally occurring, and possibly defective, mammalian polynucleotide sequence or a heterologous sequence present due to an intracellular or extracellular pathogenic infection or a disease, which polynucleotide sequence has a function that is desired to be reduced or inhibited. This target sequence may be a coding sequence, that is, it is translated to express a protein or a functional fragment thereof. Alternatively, the target sequence may be non-coding, but may have a regulatory function. One target polynucleotide sequence is a virus polynucleotide sequence necessary for replication and/or pathogenesis of the virus in an infected mammalian cell. Another embodiment of a target polynucleotide sequence is a tumor antigen or functional fragment thereof, or a non-expressed regulatory sequence of a virus-induced cancer, which sequence is required for the maintenance of the tumor in the mammal. Still another embodiment of a target polynucleotide sequence is a polynucleotide sequence of an intracellular or extracellular pathogen necessary for replication and/or pathogenesis of that pathogen in an infected mammal. Yet another embodiment of a target polynucleotide sequence is a polynucleotide sequence of an abnormal cancer-causing gene (or a non-expressed regulatory sequence) in a mammal which also possesses a normal copy of the gene or sequence. The differences between the abnormal sequence and the normal sequence are differences at the polynucleotide sequence level. Such an abnormal sequence can be, for example, a fusion of two normal genes, and the target sequence can be the sequence which spans that fusion, e.g., the BCR-abl gene sequence characteristic of certain leukemias. The term "gene" is intended to include any target sequence intended to be "silenced", whether or not transcribed and/or translated, including regulatory sequences, such as promoters.

The terms “mammal” or “mammalian” are intended to encompass their normal meaning. While the invention is most desirably intended for efficacy in humans, it may also be employed in domestic mammals such as canines, felines, and equines, as well as in mammals of particular interest, e.g., zoo animals, farmstock and the like.

5     **A.     *The Compositions of the Invention***

A composition for inhibiting the function of a target polynucleotide sequence in a mammalian cell, according to this invention, comprises an agent that provides to a mammalian cell an at least partially double-stranded RNA molecule. In general, the term “RNA” may also include RNA-DNA hybrids, except where specified otherwise,  
10     e.g., where a 2' –OH group of ribose is required for a particular linkage. The RNA molecule comprises a polynucleotide sequence of at least about 200 nucleotides in length. Importantly, this polynucleotide sequence of the RNA molecule is substantially homologous to the target polynucleotide sequence. This polynucleotide sequence also preferably contains exon sequences or portions thereof. Desirably, the  
15     polynucleotide sequence does not contain intron sequences. Preferably, the RNA molecule does not produce a functional protein, and more preferably, it is not translated. The polynucleotide sequence of the RNA molecule is preferably substantially non-homologous to any naturally occurring, normally functioning, essential mammalian polynucleotide sequence. The polynucleotide sequences  
20     described herein may employ a multitarget or polyepitope approach, e.g., encoding sequences of more than one gene of a single target pathogen or against more than one target pathogen, or other category of target desired to be silenced.

The “at least partially double stranded RNA molecule” includes an RNA polynucleotide sequence of between about 100 to 10,000 polynucleotides in length.  
25     At present the sequence is most desirably at least 200 polynucleotides in length, but it can range in one embodiment from 200 to 8000 polynucleotides in length. In another embodiment, the RNA molecule can be less than 7500 polynucleotides in length. In still another embodiment the RNA molecule can have a sequence length less than about 5000 polynucleotides. In yet another embodiment the RNA molecule can have

a sequence length less than about 2000 polynucleotides. In still another embodiment the RNA molecule can have a sequence length less than about 1000 polynucleotides. In yet another embodiment the RNA molecule can have a sequence length less than about 750 polynucleotides.

5 Minimally, to keep the RNA molecule stable, it has a minimum of 11 to 30 nucleotides involved in a double-stranded sequence, depending upon the composition of the polynucleotide sequence and a  $\Delta G$  of about -9.2 kcal/mol. As known in the art,  $\Delta G$  defines the state of minimal external energy required to keep a molecular configuration stable [see, e.g., Jaeger *et al*, Proc. Natl. Acad. Sci., USA, 20:7706-  
10 7710 (1989); and Soler and Jankowski, Math. Biosci., 2:167-190 (1991)]. Based on this minimum, preferably at least 10% of this partially double-stranded RNA molecule sequence is double-stranded. Alternatively, the double stranded portion of these RNA molecules can be at least 30% of the sequence. In another embodiment, the double stranded portion of these molecules can be at least 50% of the sequence. In still  
15 another embodiment, the double stranded portion of these molecules can be at least 70% of the sequence. In another embodiment, the double stranded portion of these molecules can be at least 90% of the sequence. In another embodiment, the entire sequence can be double stranded. Alternatively, the double-stranded portion of these molecules can occur at either or both termini, or in some middle portion of the  
20 sequence, if the molecule is linear. Similarly, the double-stranded portion can be in any location if the molecule is circular. In certain embodiments of the present invention, the double-stranded portion of the RNA molecule becomes double-stranded only when the molecule is in the mammalian cell. In still other embodiment of this invention, the partially double-stranded molecule is an RNA/DNA hybrid, for  
25 example, a single chain containing RNA and DNA, prepared *in vitro*; or a duplex of two such single chains or portions thereof. In yet another embodiment, the RNA molecule, made *in vivo* or *in vitro*, is a duplex comprised of an RNA single strand and a DNA single strand.

The partially double-stranded RNA molecule polynucleotide sequence must be  
30 substantially homologous to the target polynucleotide sequence in order to effectively

reduce or inhibit the function thereof. The necessary homology may be suitably defined by use of a computer algorithm. As known in the art, "homology" or "identity" means the degree of sequence relatedness between two polypeptide or two polynucleotide sequences as determined by the identity of the match between two lengths of such sequences. Both identity and homology can be readily calculated by methods extant in the prior art [See, e.g., COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, (1988); BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, (1993); COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, (1994); SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, (1987); and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, (1991)]. While there exist a number of methods to measure identity and homology between two polynucleotide sequences, the terms "identity", "similarity" and homology are well known to skilled artisans [H. Carillo and D. Lipton, SIAM J. Applied Math., 48:1073 (1988)]. Methods commonly employed to determine identity or homology between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and H. Carillo and D. Lipton, SIAM J. Applied Math., 48:1073 (1988). Preferred methods to determine identity or homology are designed to give the largest match between the two sequences tested. Methods to determine identity and similarity are codified in computer programs. Preferred computer program to determine identity and homology between two sequences include, but are not limited to, the algorithm BESTFIT from the GCG program package [J. Devereux et al., Nucl. Acids Res., 12(1):387 (1984)], the related MACVECTOR program (Oxford), and the FASTA (Pearson) programs. For instance, searches for sequence similarities in databases between significant naturally occurring mammalian polynucleotide sequences and target polynucleotide sequences enable the design of suitable RNA molecules desired for use in the invention. The algorithm and/or the degree of homology necessary for

any particular RNA molecule may be selected by one of skill in the art, depending on the identity of the target, and/or the closeness of homology of the target sequence to any naturally occurring mammalian sequence, which is desired to be left functioning normally after use of the methods of this invention.

5           In one preferred embodiment, the RNA polynucleotide sequence desirably has an overall homology of at least 10% to the target sequence and contains at least one segment (window) of 30 contiguous nucleotides with a homology in that window of at least 50% to a similar 30 nts region of the target sequence, using the MACVECTOR program with a default annealing temperature of 37°C. In another  
10       embodiment, the RNA polynucleotide sequence desirably has an overall homology of at least 30% to the target sequence and contains at least one window of 30 contiguous nucleotides with a homology in that window of at least 50% to a similar 30 nts region of the target sequence. In another embodiment, the RNA polynucleotide sequence desirably has an overall homology of at least 50% to the target sequence and contains  
15       at least one window of 30 contiguous nucleotides with a homology in that window of at least 50% to a similar 30 nts region of the target sequence. In another embodiment, the RNA polynucleotide sequence desirably has an overall homology of at least 70% to the target sequence and contains at least one window of 30 contiguous nucleotides with a homology in that window of at least 50% to a similar 30 nts region of the  
20       target sequence. In another embodiment, the RNA polynucleotide sequence desirably has an overall homology of at least 90% to the target sequence and contains at least one window of 30 contiguous nucleotides with a homology in that window of at least 50% to a similar 30 nts region of the target sequence.

          In still another embodiment, the RNA polynucleotide sequence desirably has  
25       an overall homology of at least 10% to the target sequence and contains at least one windows of 30 contiguous nucleotides with a homology in that window of at least 70% to a similar 30 nts region of the target sequence. In another embodiment, the RNA polynucleotide sequence desirably has an overall homology of at least 10% to the target sequence and contains at least one segment (window) of 30 contiguous

nucleotides with a homology in that window of at least 90% to a similar 30 nts region of the target sequence.

In yet another embodiment, the RNA polynucleotide sequence desirably has an overall homology of at least 10% to the target sequence and contains at least two  
5 windows of 30 contiguous nucleotides with a homology in the windows of at least 50% to similar 30 nts regions of the target sequence. Other embodiments of this formula can be developed by one of skill in the art.

In a second preferred embodiment, the RNA polynucleotide sequence desirably has an overall homology of at least 10% to the target sequence and contains  
10 at least one segment (window) of 5 contiguous nucleotides with absolute homology in that window to a 5 nts region of the target sequence, using the MACVECTOR program with a default annealing temperature of 37°C. In another variant of this embodiment, the RNA polynucleotide sequence desirably has an overall homology of at least 30% to the target sequence and contains at least one window of 5 contiguous  
15 nucleotides with absolute homology to a 5 nts region of the target sequence. In another embodiment, the RNA polynucleotide sequence desirably has an overall homology of at least 50% to the target sequence and contains the above described 5 nt absolutely homologous window. Other variants of this embodiment can be developed by one of skill in the art.

20 The presence of the windows referred to in the formulae above permits the overall homology of the remainder of the sequence to be low; however it is anticipated that a low overall homology is likely to affect the dosage of the therapeutic compositions described below adversely. An increase in the number of such windows in the RNA polynucleotide sequence is likely to permit the overall homology of the  
25 rest of the sequence to be low, but not affect the dosage

It should be understood that selection of the necessary homology, selection of the defaults for the program and selection of the program employed to calculate homology is within the skill of the art, given the teachings of this specification and the knowledge extant in the scientific literature.

The RNA molecule polynucleotide sequence is also desirably substantially non-homologous to any naturally occurring, normally functioning, and essential mammalian polynucleotide sequence, so that the RNA molecule polynucleotide sequence does not adversely affect the function of any essential naturally occurring mammalian polynucleotide sequence, when used in the methods of this invention. Such naturally occurring functional mammalian polynucleotide sequences include mammalian sequences that encode desired proteins, as well as mammalian sequences that are non-coding, but that provide for essential regulatory sequences in a healthy mammal. Essentially, the RNA molecule useful in this invention must be sufficiently distinct in sequence from any mammalian polynucleotide sequence for which the function is intended to be undisturbed after any of the methods of this invention is performed. As described for determining the homology to the target sequence above, one of skill in the art may have resort to the above-identified computer algorithms to define the essential lack of homology between the RNA molecule polynucleotide sequence and the normal mammalian sequences. Thus, in one exemplary embodiment, the homology between the RNA polynucleotide and the selected normal sequence is less than the homologies of the formulae described above. More preferably, there is almost no homology at all between the RNA polynucleotide and any normal mammalian sequence. It should be understood that selection of the necessary homology is within the skill of the art, given the teachings of this specification and the knowledge extant in the scientific literature.

Finally, yet another desirable attribute of the RNA molecule of the composition of the present invention is that it does not produce a functional protein, or alternatively, is not translated. The RNA molecule or the delivery agent can be engineered in a variety of known ways, so as to optionally not express a functional protein or to optionally not interact with cellular factors involved in translation. Thus, for example, the agent, whether it be a synthesized RNA molecule or an agent which becomes an RNA molecule *in vivo*, lacks a poly-adenylation sequence. Similarly, the agent can lack a Kozak region necessary for protein translation. In another embodiment, the RNA molecule can also lack the native initiating methionine codon.



In still another embodiment, the RNA molecule polynucleotide sequence lacks a cap structure. In yet a further embodiment, the RNA molecule has no signals for protein synthesis. In still another embodiment, the RNA molecule contains no coding sequence or a functionally inoperative coding sequence. In still another embodiment, the RNA sequence can be punctuated with intronic sequences. In yet a further embodiment, a hairpin sequence can be placed before the native initiation codon, if present. In still another embodiment, the RNA molecule can be an RNA/DNA hybrid as described above. All such embodiments can be designed by resort to the known teachings of, e.g., such texts as cited below.

10           The following are various specific embodiments that may be used to achieve polynucleotide inhibition as described herein. It should be recognized that the various RNA (and RNA/DNA hybrid) structures described below may be used singly or in any combination of two or more, e.g., a lariat (sense or antisense) and/or a complementary circular and/or linear molecule. The antisense lariat or circle structures may also be used alone. Furthermore, these structures may include regions of self complementarity (e.g., tandem sense and antisense sequences) as well as additional antisense sequences relative to a desired target. Throughout this document the term "antisense" is used to mean complementary to and capable of hybridizing with any mRNA.

20           In one embodiment, polynucleotides in the form of "lariats" may be utilized. Lariats contain a 2'-5' phosphodiester linkage as opposed to the usual 3'-5' linkage. Such structures are formed in splicing reactions catalyzed by spliceosomes and self-cleaving ribozymes. These structures are either intermediates or by-products of splicing reactions. They can be prepared *in vivo* through expression (transcription) in a cell or prepared *in vitro*. Lariats form when a free 5' phosphoryl group of either a ribose or deoxyribose becomes linked to the 2' -OH of a ribose in a loop back fashion. The lariats may contain 10 or more nucleotides in the loop or may be a complete circle, with the loop back linkage in each case being 2'-5'. A lariat linking the terminal nucleotides produces a circle-like structure. The loops and/or the stem can contain either the sense and the antisense sequences in tandem in a single molecule, or

each single lariat contains either a sense or an antisense sequence. The lariats that contain sense and antisense in separate molecules may be administered together as a double-stranded form or the antisense lariat may be used singly to form a double strand with the mRNA in the cell. Lariats may be RNA or a DNA hybrid, with the 2'-5' linkage effected through the 2' -OH of the RNA portion of the hybrid [Rees C and Song Q. Nucl. Acid Res., 27, 2672-2681 (1999); Dame E *et al*, Biochemistry, 38, 3157-3167, 1999; Clement J.Q. *et al*, RNA, 5, 206-220, 1999; Block T and Hill J. J. Neurovirol., 3, 313-321, 1997; Schindewolf CA and Domdey H., Nucl. Acid Res., 23, 1133-1139 (1995)].

10 In another embodiment, a circular RNA (or circular RNA-DNA hybrid) can be generated through a 2'-5' or a 3'-5' linkage of the termini. These may be generated enzymatically through RNA ligase reactions using a splinter to bring the ends in proximity *in vitro*, or through the use of self splicing ribozymes (*in vivo* and *in vitro*). The desired inhibition may be achieved by providing one or more RNA circles, made  
15 *in vitro* or expressed *in vivo*, including single circles with or without self complementarity, as well as double stranded circular RNA (both sense and antisense strands relative to the target polynucleotide), or two circles of single-stranded RNA which have regions of complementarity to each other as well as one having complementarity to a target.

20 Another embodiment utilizes single RNA (or RNA-DNA hybrid) antisense circles (circular RNA without self complementarity which is complementary to the target mRNA). Still another embodiment utilizes RNA-DNA circles or a circular DNA molecule complementary to a target mRNA molecule. Single circles with tandem sense and antisense sequences (in any order) which have complementarity to a  
25 target message may be used as the composition which inhibits the function of the target sequence. It may be preferred to use circular molecules having such self-complementary sequences which may form rod-like sections, as well as additional antisense sequences to the target [Schindewolf CA and Domdey H. Nucl. Acid Res., 23, 1133-1139 (1995); Rees C and Song Q., Nucl. Acid Res., 27, 2672-2681 (1999);  
30 Block T and Hill J., J. Neurovirol., 3, 313-321(1997)].

In yet a further embodiment, the composition which inhibits the target sequence is a capped linear RNA. Whether the dsRNA is formed *in vitro* or *in vivo*, either one or both strands may be capped. In circumstances where cytoplasmic expression would not ordinarily result in capping of the RNA molecule, capping may be accomplished by various means including use of a capping enzyme, such as a vaccinia capping enzyme or an alphavirus capping enzyme. A capped antisense molecule may be used to achieve the desired post transcriptional silencing of the target gene. Capped RNA may be prepared *in vitro* or *in vivo*. RNA made in the nucleus by RNA polII ordinarily is capped. Cytoplasmically expressed RNA may or may not be capped. Capping can be achieved by expressing capping enzymes of cytoplasmic viruses. Either both capped or one capped and one uncapped or both uncapped RNA or RNA-DNA hybrid sequences may be used in these compositions. Capped or uncapped antisense molecule may be used, singly or in any combination with polynucleotide structures described herein.

The RNA molecule according to this invention may be delivered to the mammalian or extracellular pathogen present in the mammalian cell in the composition as an RNA molecule or partially double stranded RNA sequence, or RNA/DNA hybrid, which was made *in vitro* by conventional enzymatic synthetic methods using, for example, the bacteriophage T7, T3 or SP6 RNA polymerases according to the conventional methods described by such texts as the Promega Protocols and Applications Guide, (3rd ed. 1996), eds. Doyle, ISBN No. 1-882274-57-1.

Alternatively these molecules may be made by chemical synthetic methods *in vitro* [see, e.g., Q. Xu *et al*, Nucl. Acids Res., 24(18):3643-4 (Sept. 1996); N. Naryshkin *et al*, Bioorg. Khim., 22(9):691-8 (Sept. 1996); J. A. Grasby *et al*, Nucl. Acids Res., 21(19):4444-50 (Sept 1993); C. Chaix *et al*, Nucl. Acids Res., 17(18):7381-93 (1989); S.H. Chou *et al*, Biochem., 28(6):2422-35 (Mar. 1989); O. Odai *et al*, Nucl. Acids Symp. Ser., 21:105-6 (1989); N.A. Naryshkin *et al*, Bioorg. Khim., 22(9):691-8 (Sept. 1996); S. Sun *et al*, RNA, 3(11):1352-1363 (Nov. 1997); X. Zhang *et al*, Nucl. Acids Res., 25(20):3980-3 (Oct. 1997); S. M. Grvaznov *et al*,

Nucl. Acids Res., 26(18):4160-7 (Sept. 1998); M. Kadokura *et al*, Nucl. Acids Symp Ser, 37:77-8 (1997); A. Davison *et al*, Biomed. Pept. Proteins. Nucl. Acids, 2(1):1-6 (1996); and A. V. Mudrakovskaia *et al*, Bioorg. Khim., 17(6):819-22 (Jun. 1991)].

Still alternatively, the RNA molecule of this invention can be made in a  
5 recombinant microorganism, e.g., bacteria and yeast or in a recombinant host cell,  
e.g., mammalian cells, and isolated from the cultures thereof by conventional  
techniques. See, e.g., the techniques described in Sambrook et al, MOLECULAR  
CLONING, A LABORATORY MANUAL, 2nd Ed.; Cold Spring Harbor Laboratory  
Press, Cold Spring Harbor, New York, 1989, which is exemplary of laboratory  
10 manuals that detail these techniques, and the techniques described in US Patent Nos.  
5,824,538; 5,877,159 and 65,643,771, incorporated herein by reference.

Such RNA molecules prepared or synthesized *in vitro* may be directly  
delivered to the mammalian cell or to the mammal as they are made *in vitro*. The  
references above provide one of skill in the art with the techniques necessary to  
15 produce any of the following specific embodiments, given the teachings provided  
herein. Therefore, in one embodiment, the "agent" of the composition is a duplex  
(i.e., it is made up of two strands), either complete or partially double stranded RNA.  
In another embodiment, the agent is a single stranded RNA sense strand. In another  
embodiment, the agent of the composition is a single stranded RNA anti-sense strand.  
20 Preferably the single stranded RNA sense or anti-sense strand forms a hairpin at one  
or both termini. Desirably, the single stranded RNA sense or anti-sense strand forms  
a hairpin at some intermediate portion between the termini. Such a single stranded  
RNA sense or anti-sense strand may also be designed to fold back upon itself to  
become partially double stranded *in vitro* or *in vivo*. Yet another embodiment of an  
25 extant RNA molecule as the effective agent used in the compositions is a single  
stranded RNA sequence comprising both a sense polynucleotide sequence and an anti-  
sense polynucleotide sequence, optionally separated by a non-base paired  
polynucleotide sequence. Preferably, this single stranded RNA sequence has the  
ability to become double-stranded once it is in the cell, or *in vitro* during its synthesis.  
30 Still another embodiment of this invention is an RNA/DNA hybrid as described above.

Still another embodiment of the synthetic RNA molecule is a circular RNA molecule that optionally forms a rod structure [see, e.g., K-S. Wang *et al*, Nature, **323**:508-514 (1986)] or is partially double-stranded, and can be prepared according to the techniques described in S. Wang *et al*, Nucl. Acids Res., **22**(12):2326-33 (June 1994);  
5 Y. Matsumoto *et al*, Proc. Natl. Acad. Sci. USA, **87**(19):7628-32 (Oct. 1990); Proc. Natl. Acad. Sci. USA, **91**(8):3117-21 (Apr. 1994); M. Tsagris *et al*, Nucl. Acids Res., **19**(7):1605-12 (Apr. 1991); S. Braun *et al*, Nucl. Acids Res., **24**(21):4152-7 (Nov. 1996); Z. Pasman *et al*, RNA, **2**(6):603-10 (Jun. 1996); P. G. Zaphiropoulos, Proc. Natl. Acad. Sci. USA, **93**(13):6536-41 (Jun. 1996); D. Beaudry *et al*, Nucl.  
10 Acids Res., **23**(15):3064-6 (Aug. 1995), all incorporated herein by reference. Still another agent is a double-stranded molecule comprised of RNA and DNA present on separate strands, or interspersed on the same strand.

Alternatively, the RNA molecule may be formed *in vivo* and thus delivered by a "delivery agent" which generates such a partially double-stranded RNA molecule *in*  
15 *vivo* after delivery of the agent to the mammalian cell or to the mammal. Thus, the agent which forms the composition of this invention is, in one embodiment, a double stranded DNA molecule "encoding" one of the above-described RNA molecules. The DNA agent provides the nucleotide sequence which is transcribed within the cell to become a double stranded RNA. In another embodiment, the DNA sequence  
20 provides a deoxyribonucleotide sequence which within the cell is transcribed into the above-described single stranded RNA sense or anti-sense strand, which optionally forms a hairpin at one or both termini or folds back upon itself to become partially double stranded. The DNA molecule which is the delivery agent of the composition can provide a single stranded RNA sequence comprising both a sense polynucleotide  
25 sequence and an anti-sense polynucleotide sequence, optionally separated by a non-base paired polynucleotide sequence, and wherein the single stranded RNA sequence has the ability to become double-stranded. Alternatively, the DNA molecule which is the delivery agent provides for the transcription of the above-described circular RNA molecule that optionally forms a rod structure or partial double strand *in vivo*. The  
30 DNA molecule may also provide for the *in vivo* production of an RNA/DNA hybrid as

described above, or a duplex containing one RNA strand and one DNA strand. These various DNA molecules may be designed by resort to conventional techniques such as those described in Sambrook, cited above or in the Promega reference, cited above.

A latter delivery agent of the present invention, which enables the formation in  
5 the mammalian cell of any of the above-described RNA molecules, can be a DNA single stranded or double stranded plasmid or vector. Expression vectors designed to produce RNAs as described herein *in vitro* or *in vivo* may containing sequences under the control of any RNA polymerase, including mitochondrial RNA polymerase, RNA polI, RNA polII, and RNA polIII. These vectors can be used to transcribe the desired  
10 RNA molecule in the cell according to this invention. Vectors may be desirably designed to utilize an endogenous mitochondrial RNA polymerase (e.g., human mitochondrial RNA polymerase, in which case such vectors may utilize the corresponding human mitochondrial promoter). Mitochondrial polymerases may be used to generate capped (through expression of a capping enzyme) or uncapped  
15 messages *in vivo*. RNA pol I, RNA pol II, and RNA pol III transcripts may also be generated *in vivo*. Such RNAs may be capped or not, and if desired, cytoplasmic capping may be accomplished by various means including use of a capping enzyme such as a vaccinia capping enzyme or an alphavirus capping enzyme. The DNA vector is designed to contain one of the promoters or multiple promoters in  
20 combination (mitochondrial, RNA polI, II, or polIII, or viral, bacterial or bacteriophage promoters along with the cognate polymerases). Preferably, where the promoter is RNA pol II, the sequence encoding the RNA molecule has an open reading frame greater than about 300 nts to avoid degradation in the nucleus. Such plasmids or vectors can include plasmid sequences from bacteria, viruses or phages.  
25 Such vectors include chromosomal, episomal and virus-derived vectors e.g., vectors derived from bacterial plasmids, bacteriophages, yeast episomes, yeast chromosomal elements, and viruses, vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, cosmids and phagemids. Thus, one exemplary vector is a single or double-stranded phage vector. Another  
30 exemplary vector is a single or double-stranded RNA or DNA viral vector. Such

vectors may be introduced into cells as polynucleotides, preferably DNA, by well known techniques for introducing DNA and RNA into cells. The vectors, in the case of phage and viral vectors may also be and preferably are introduced into cells as packaged or encapsidated virus by well known techniques for infection and  
5 transduction. Viral vectors may be replication competent or replication defective. In the latter case, viral propagation generally occurs only in complementing host cells.

In another embodiment the delivery agent comprises more than a single DNA or RNA plasmid or vector. As one example, a first DNA plasmid can provide a single stranded RNA sense polynucleotide sequence as described above, and a second DNA  
10 plasmid can provide a single stranded RNA anti-sense polynucleotide sequence as described above, wherein the sense and anti-sense RNA sequences have the ability to base-pair and become double-stranded. Such plasmid(s) can comprise other conventional plasmid sequences, e.g., bacterial sequences such as the well-known sequences used to construct plasmids and vectors for recombinant expression of a  
15 protein. However, it is desirable that the sequences which enable protein expression, e.g., Kozak regions, etc., are not included in these plasmid structures.

The vectors designed to produce dsRNAs of the invention may desirably be designed to generate two or more, including a number of different dsRNAs homologous and complementary to a target sequence. This approach is desirable in  
20 that a single vector may produce many, independently operative dsRNAs rather than a single dsRNA molecule from a single transcription unit and by producing a multiplicity of different dsRNAs, it is possible to self select for optimum effectiveness. Various means may be employed to achieve this, including autocatalytic sequences as well as sequences for cleavage to create random and/or predetermined splice sites.

25 Other delivery agents for providing the information necessary for formation of the above-described desired RNA molecules in the mammalian cell include live, attenuated or killed, inactivated recombinant bacteria which are designed to contain the sequences necessary for the required RNA molecules of this invention. Such recombinant bacterial cells, fungal cells and the like can be prepared by using  
30 conventional techniques such as described in US Patent Nos. 5,824,538; 5,877,159

and 65,643,771, incorporated herein by reference. Microorganisms useful in preparing these delivery agents include those listed in the above cited reference, including, without limitation, *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various species of *Pseudomonas*, *Streptomyces*, and

5 *Staphylococcus*.

Still other delivery agents for providing the information necessary for formation of the desired, above-described RNA molecules in the mammalian cell include live, attenuated or killed, inactivated viruses, and particularly recombinant viruses carrying the required RNA polynucleotide sequence discussed above. Such

10 viruses may be designed similarly to recombinant viruses presently used to deliver genes to cells for gene therapy and the like, but preferably do not have the ability to express a protein or functional fragment of a protein. Among useful viruses or viral sequences which may be manipulated to provide the required RNA molecule to the mammalian cell *in vivo* are, without limitation, alphavirus, adenovirus, adeno-

15 associated virus, baculoviruses, delta virus, pox viruses, hepatitis viruses, herpes viruses, papova viruses (such as SV40), poliovirus, pseudorabies viruses, retroviruses, vaccinia viruses, positive and negative stranded RNA viruses, viroids, and virusoids, or portions thereof. These various viral delivery agents may be designed by applying conventional techniques such as described in M. Di Nocola *et al*, Cancer Gene Ther.,

20 5(6):350-6 (1998), among others, with the teachings of the present invention.

Another delivery agent for providing the information necessary for formation of the desired, above-described RNA molecules in the mammalian cell include live, attenuated or killed, inactivated donor cells which have been transfected or infected *in vitro* with a synthetic RNA molecule or a DNA delivery molecule or a delivery

25 recombinant virus as described above. These donor cells may then be administered to the mammal, as described in detail below, to stimulate the mechanism in the mammal which mediates this inhibitory effect. These donor cells are desirably mammalian cells, such as C127, 3T3, CHO, HeLa, human kidney 293, BHK cell lines, and COS-7 cells, and preferably are of the same mammalian species as the mammalian recipient. Such

30 donor cells can be made using techniques similar to those described in, e.g., Emerich



*et al*, J. Neurosci., 16: 5168-81 (1996). Even more preferred, the donor cells may be harvested from the specific mammal to be treated and made into donor cells by *ex vivo* manipulation, akin to adoptive transfer techniques, such as those described in D. B. Kohn *et al*, Nature Med., 4(7):775-80 (1998). Donor cells may also be from non-mammalian species, if desired.

Finally, the composition of this invention can also include one or more of the selected agents which are described above. The composition can contain a mixture of synthetic RNA molecules described above, synthetic DNA delivery molecules described above, and any of the other delivery agents described above, such as recombinant bacteria, cells, and viruses. The identity of the composition mixture may be readily selected by one of skill in the art.

***B. Pharmaceutical (Therapeutic or Prophylactic) Compositions of the Invention***

The compositions of this invention for pharmaceutical use desirably contain the synthetic RNA molecule as described above or the agent which provides that RNA molecule to the mammalian cell *in vivo* in a pharmaceutically acceptable carrier, with additional optional components for pharmaceutical delivery. The specific formulation of the pharmaceutical composition depends upon the form of the agent delivering the RNA molecule.

Suitable pharmaceutically acceptable carriers facilitate administration of the polynucleotide compositions of this invention, but are physiologically inert and/or nonharmful. Carriers may be selected by one of skill in the art. Such carriers include but are not limited to, sterile saline, phosphate, buffered saline, dextrose, sterilized water, glycerol, ethanol, lactose, sucrose, calcium phosphate, gelatin, dextran, agar, pectin, peanut oil, olive oil, sesame oil, and water and combinations thereof.

Additionally, the carrier or diluent may include a time delay material, such as glycerol monostearate or glycerol distearate alone or with a wax. In addition, slow release polymer formulations can be used. The formulation should suit not only the form of

the delivery agent, but also the mode of administration. Selection of an appropriate carrier in accordance with the mode of administration is routinely performed by those skilled in the art.

Where the composition contains the synthetic RNA molecule or where the agent is another polynucleotide, such as, a DNA molecule, plasmid, viral vector, or recombinant virus, or multiple copies of the polynucleotide or different polynucleotides, etc., as described above, the composition may desirably be formulated as "naked" polynucleotide with only a carrier. Alternatively, such compositions desirably contain optional polynucleotide facilitating agents or "co-agents", such as a local anaesthetic, a peptide, a lipid including cationic lipids, a liposome or lipidic particle, a polycation such as polylysine, a branched, three-dimensional polycation such as a dendrimer, a carbohydrate, a cationic amphiphile, a detergent, a benzylammonium surfactant, or another compound that facilitates polynucleotide transfer to cells. Non-exclusive examples of such facilitating agents or co-agents useful in this invention are described in U. S. Patent Nos. 5,593,972; 5,703,055; 5,739,118; 5,837,533 and International Patent Application No. WO96/10038, published April 4, 1996; and International Patent Application No. WO94/16737, published August 8, 1994, which are each incorporated herein by reference.

When the facilitating agent used is a local anesthetic, preferably bupivacaine, an amount of from about 0.1 weight percent to about 1.0 weight percent based on the total weight of the polynucleotide composition is preferred. See, also, International Patent Application No. PCT/US98/22841, which teaches the incorporation of benzylammonium surfactants as co-agents, administered in an amount of between about 0.001-0.03 weight %, the teaching of which is hereby incorporated by reference.

Where the delivery agent of the composition is other than a polynucleotide composition, e.g., is a transfected donor cell or a bacterium as described above, the composition may also contain other additional agents, such as those discussed in US Patents No. 5,824,538; 5,643,771; 5,877,159, incorporated herein by reference.

Still additional components that may be present in any of the compositions are, adjuvants, preservatives, chemical stabilizers, or other antigenic proteins. Typically, stabilizers, adjuvants, and preservatives are optimized to determine the best formulation for efficacy in the target human or animal. Suitable exemplary

5 preservatives include chlorobutanol, potassium sorbate, sorbic acid, sulfur dioxide, propyl gallate, the parabens, ethyl vanillin, glycerin, phenol, and parachlorophenol. Suitable stabilizing ingredients which may be used include, for example, casamino acids, sucrose, gelatin, phenol red, N-Z amine, monopotassium diphosphate, lactose, lactalbumin hydrolysate, and dried milk. A conventional adjuvant is used to attract

10 leukocytes or enhance an immune response. Such adjuvants include, among others, Ribi, mineral oil and water, aluminum hydroxide, Amphigen, Avridine, L121/squalene, D-lactide-poly(lactide)/glycoside, pluronic polyols, muramyl dipeptide, killed *Bordetella*, and saponins, such as Quil A.

In addition, other agents which may function as transfecting agents and/or

15 replicating agents and/or inflammatory agents and which may be co-administered with the composition of this invention, include growth factors, cytokines and lymphokines such as alpha-interferon, gamma-interferon, platelet derived growth factor (PDGF), colony stimulating factors, such as G-CSF, GM-CSF, tumor necrosis factor (TNF), epidermal growth factor (EGF), and the interleukins, such as IL-1, IL-2, IL-4, IL-6,

20 IL-8, IL-10 and IL-12. Further, fibroblast growth factor, surface active agents such as immune-stimulating complexes (ISCOMS), Freund's incomplete adjuvant, LPS analog including monophosphoryl Lipid A (MPL), muramyl peptides, quinone analogs and vesicular complexes such as squalene and squalene, and hyaluronic acid may also be used administered in conjunction with the compositions of the invention.

25 The pharmaceutical compositions may also contain other additives suitable for the selected mode of administration of the composition. Thus, these compositions can contain additives suitable for administration via any conventional route of administration, including without limitation, parenteral administration, intraperitoneal administration, intravenous administration, intramuscular administration,

30 subcutaneous administration, intradermal administration, oral administration, topical

administration, intranasal administration, intra-pulmonary administration, rectal administration, vaginal administration, and the like. All such routes are suitable for administration of these compositions, and may be selected depending on the agent used, patient and condition treated, and similar factors by an attending physician.

5           The composition of the invention may also involve lyophilized polynucleotides, which can be used with other pharmaceutically acceptable excipients for developing powder, liquid or suspension dosage forms, including those for intranasal or pulmonary applications. See, e.g., Remington: The Science and Practice of Pharmacy, Vol. 2, 19<sup>th</sup> edition (1995), e.g., Chapter 95 Aerosols; and International Patent  
10   Application No. PCT/US99/05547, the teachings of which are hereby incorporated by reference. Routes of administration for these compositions may be combined, if desired, or adjusted.

          In some preferred embodiments, the pharmaceutical compositions of the invention are prepared for administration to mammalian subjects in the form of, for  
15   example, liquids, powders, aerosols, tablets, capsules, enteric coated tablets or capsules, or suppositories.

          The compositions of the present invention, when used as pharmaceutical compositions, can comprise about 1 ng to about 20 mgs of polynucleotide molecules as the delivery agent of the compositions, e.g., the synthetic RNA molecules or the  
20   delivery agents which can be DNA molecules, plasmids, viral vectors, recombinant viruses, and mixtures thereof. In some preferred embodiments, the compositions contain about 10 ng to about 10 mgs of polynucleotide sequences. In other embodiments, the pharmaceutical compositions contain about 0.1 to about 500  $\mu$ g polynucleotide sequences. In some preferred embodiments, the compositions contain  
25   about 1 to about 350  $\mu$ g polynucleotide sequences. In still other preferred embodiments, the pharmaceutical compositions contain about 25 to about 250  $\mu$ g of the polynucleotide sequences. In some preferred embodiments, the vaccines and therapeutics contain about 100  $\mu$ g of the polynucleotide sequences.

          The compositions of the present invention in which the delivery agents are  
30   donor cells or bacterium can be delivered in dosages of between about 1 cell to about

10<sup>7</sup> cells/dose. Similarly, where the delivery agent is a live recombinant virus, a suitable vector-based composition contains between 1x10<sup>2</sup> pfu to 1x10<sup>12</sup> pfu per dose.

Given the teachings of this invention, and the observed capacity of the inhibitory effect of the methods and compositions of this invention to be propagated to more cells than the cells transfected or infected with the composition of this invention, it is likely that suitable dosage adjustments can be made downwards from the above-noted dosages. Thus, the above dosage ranges are guidelines only. In general, the pharmaceutical compositions are administered in an amount effective to inhibit or reduce the function of the target polynucleotide sequence for treatment or prophylaxis of the diseases, disorders or infections for which such target functions are necessary for further propagation of the disease or causative agent of the disease. The amount of the pharmaceutical composition in a dosage unit employed is determined empirically, based on the response of cells *in vitro* and response of experimental animals to the compositions of this invention. Optimum dosage is determined by standard methods for each treatment modality and indication. Thus the dose, timing, route of administration, and need for readministration of these compositions may be determined by one of skill in the art, taking into account the condition being treated, its severity, complicating conditions, and such factors as the age, and physical condition of the mammalian subject, the employment of other active compounds, and the like.

### ***C. Therapeutic and Prophylactic Methods of the Invention***

The methods of this invention can employ the compositions described in detail above, and possibly other polynucleotide sequences currently used in the art (e.g., polynucleotide molecules which do encode proteins, whether functional or non-functional, or known RNA catalytic sequences, such as ribozymes) which can provide partially double stranded RNA molecules to a mammalian cell. It is anticipated, however, that the efficiency of these methods is enhanced by the use of RNA molecules which do not produce protein. These methods reduce or inhibit the function of a target polynucleotide sequence(s) in a mammal or in the cell of a

mammal. The compositions, pharmaceutical compositions, dosages and modes of administration described above are particularly desirable for the treatment of a variety of disorders that plague mammals, including infections by heterologous pathogenic organisms, either extracellular or intracellular pathogens. Additionally, the  
5 compositions of this invention are useful in preventing infection of a mammal with a pathogen, or preventing the occurrence of disorders caused by reactivation of a latent pathogen. These compositions are also useful for the treatment of pathogenically-induced cancers.

One embodiment of a method of this invention is a method for treating a viral  
10 infection in a mammal. Particularly suitable for such treatment area DNA viruses or viruses that have an intermediary DNA stages. Among such viruses are included, without limitation, viruses of the species Retrovirus, Herpesvirus, Hepadenovirus, Poxvirus, Parvovirus, Papillomavirus, and Papovavirus. Specifically some of the more desirable viruses to treat with this method include, without limitation, HIV, HBV,  
15 HSV, CMV, HPV, HTLV and EBV. The agent used in this method provides to the cell of the mammal an at least partially double stranded RNA molecule as described above, which is substantially homologous to a target polynucleotide which is a virus polynucleotide sequence necessary for replication and/or pathogenesis of the virus in an infected mammalian cell. Among such target polynucleotide sequences are protein-  
20 encoding sequences for proteins necessary for the propagation of the virus, e.g., the HIV gag, env and pol genes, the HPV6 L1 and E2 genes, the HPV11 L1 and E2 genes, the HPV16 E6 and E7 genes, the HPV18 E6 and E7 genes, the HBV surface antigens, the HBV core antigen, HBV reverse transcriptase, the HSV gD gene, the HSVvp16 gene, the HSV gC, gH, gL and gB genes, the HSV ICP0, ICP4 and ICP6  
25 genes, Varicella zoster gB, gC and GH genes, and the BCR-abl chromosomal sequences, and non-coding viral polynucleotide sequences which provide regulatory functions necessary for transfer of the infection from cell to cell, e.g., the HIV LTR, and other viral promoter sequences, such as HSV vp16 promoter, HSV-ICP0 promoter, HSV- ICP4, ICP6 and gD promoters, the HBV surface antigen promoter,  
30 the HBV pre-genomic promoter, among others. As described above, the composition

is administered with an polynucleotide uptake enhancer or facilitator and an optional pharmaceutically acceptable carrier. The amount or dosage which is administered to the mammal is effective to reduce or inhibit the function of the viral sequence in the cells of the mammal.

5           While not wishing to be bound by theory, once the RNA molecule is delivered to or produced in a cell infected by the virus, the exogenous RNA molecule reduces or inhibits (i.e. turns off) the homologous viral sequence and is itself inhibited, so that the function of the viral sequence is reduced or inhibited. As demonstrated in the examples below, the inhibition of function effect is transferred from the mammalian  
10   cell which receives the exogenous RNA molecule to other mammalian cells in the subject which have not directly been provided with the exogenous RNA molecule. It is presently theorized that this results occurs on the level of RNA degradation.

          Thus, this method can be used to treat mammalian subjects already infected with a virus, such as HIV, in order to shut down or inhibit a viral gene function  
15   essential to virus replication and/or pathogenesis, such as HIV gag. Alternatively, this method can be employed to inhibit the functions of viruses which exist in mammals as latent viruses, e.g., *Varicella zoster* virus, and are the causative agents of the disease known as shingles. Similarly, diseases such as atherosclerosis, ulcers, chronic fatigue syndrome, and autoimmune disorders, recurrences of HSV-1 and HSV-2, HPV  
20   persistent infection, e.g., genital warts, and chronic HBV infection among others, which have been shown to be caused, at least in part, by viruses, bacteria or another pathogen, can be treated according to this method by targeting certain viral polynucleotide sequences essential to viral replication and/or pathogenesis in the mammalian subject.

25           In still another embodiment of this invention, the compositions described above can be employed in a method to prevent viral infection in a mammal. When the method described above, i.e., administering a composition described above in an amount effective to reduce or inhibit the function of the essential target viral polynucleotide sequence to a mammal, is administered prior to exposure of the  
30   mammal to the virus, it is expected that the exogenous RNA molecule remains in the

mammal and work to inhibit any homologous viral sequence which presents itself to the mammal thereafter. Thus, the compositions of the present invention may be used to inhibit or reduce the function of a viral polynucleotide sequence for vaccine use.

Still an analogous embodiment of the above "anti-viral" methods of the invention includes a method for treatment or prophylaxis of a virally induced cancer in a mammal. Such cancers include HPV E6/E7 virus-induced cervical carcinoma, HTLV-induced cancer, and EBV induced cancers, such as Burkitts lymphoma, among others. This method is accomplished by administering to the mammal a composition as described above in which the target polynucleotide is a sequence encoding a tumor antigen or functional fragment thereof, or a non-expressed regulatory sequence, which antigen or sequence function is required for the maintenance of the tumor in the mammal. Among such sequences are included, without limitation, HPV16 E6 and E7 sequences and HPV 18 E6 and E7 sequences. Others may readily be selected by one of skill in the art. The composition is administered in an amount effective to reduce or inhibit the function of the antigen in the mammal, and preferably employs the composition components, dosages and routes of administration as described above. The molecular mechanism underlying this method is the same as that described above.

In another embodiment of the invention, the compositions of this invention can be employed in a method for the treatment or prophylaxis of infection of a mammal by a non-viral pathogen, either intracellular or extracellular. As used herein, the term "intracellular pathogen" is meant to refer to a virus, bacteria, protozoan or other pathogenic organism that, for at least part of its reproductive or life cycle, exists within a host cell and therein produces or causes to be produced, pathogenic proteins. Intracellular pathogens which infect cells which include a stage in the life cycle where they are intracellular pathogens include, without limitation, Listeria, Chlamydia, Leishmania, Brucella, Mycobacteria, Shigella, and as well as Plasmodia, e.g., the causative agent of malaria, *P. falciparum*. Extracellular pathogens are those which replicate and/or propagate outside of the mammalian cell, e.g., Gonorrhoeae, and Borrellia, among others. According to this embodiment, such infection by an pathogen may be treated or possibly prevented by administering to a mammalian



subject, either already infected or anticipating exposure to the pathogen, with a composition as described above with an optional second agent that facilitates polynucleotide uptake in a cell, in a pharmaceutically acceptable carrier. In this case, the RNA molecule of the composition has a polynucleotide sequence which is

5 substantially homologous to a target polynucleotide sequence of the pathogen that is necessary for replication and/or pathogenesis of the pathogen in an infected mammal or mammalian cell. As above, the amount of the composition administered is an amount effective to reduce or inhibit the function of the pathogenic sequence in the mammal. The dosages, timing, routes of administration and the like are as described

10 above.

One of skill in the art, given this disclosure can readily select viral families and genera, or pathogens including prokaryotic and eukaryotic protozoan pathogens as well as multicellular parasites, for which therapeutic or prophylactic compositions according to the present invention can be made. See, e.g., the tables of such

15 pathogens in general immunology texts and in U. S. Patent No. 5,593,972, incorporated by reference herein.

The compositions of this invention and possibly protein-encoding molecules of the prior art may also be employed in another novel method of this invention. Such compositions are also useful in the treatment of certain non-pathogenic diseases or

20 disorders of mammals, such as certain cancers or inherited disorders. Among conditions particularly susceptible to treatment or prophylaxis according to this invention are those conditions which are characterized by the presence of an aberrant mammalian polynucleotide sequence, the function of which is necessary to the initiation or progression of the disorder, but can be inhibited without causing harm or

25 otherwise unduly adversely impacting the health of the mammal. In other words, a characteristic of a disorder suitable for this treatment is that the mammal can survive without the function of the gene, or can survive if the function of the gene was substantially reduced. In such cases, the function of the aberrant or abnormal polynucleotide sequence can be replaced exogenously by therapy. In another case, the

30 disease can be caused by the presence or function of an abnormal polynucleotide

sequence or gene in a mammal, where the mammal also possesses a normal copy of the polynucleotide sequence or gene, and wherein the differences between the abnormal gene and the normal gene are differences in nucleotide sequence. In such cases, inhibition of the function of the abnormal polynucleotide sequence by the  
5 method of this invention is likely to permit the normal polynucleotide sequence to function, without exogenous treatment.

Thus, in one embodiment, a method of treatment or prophylaxis of a cancer in a mammal involves administering to the mammal a composition of this invention in which the target polynucleotide sequence is an abnormal cancer-causing  
10 polynucleotide sequence or gene in a mammal. The composition of this invention is administered in an amount effective to reduce or inhibit the function of the abnormal sequence in the mammal. As described above, the composition can contain an optional second agent that facilitates polynucleotide uptake in a cell, and a pharmaceutically acceptable carrier, and be administered in dosages, regimens and by  
15 routes as described above.

Mammalian cancers which are characterized by the presence of abnormal and normal polynucleotide sequences include chronic myelogenous leukemia (CML) and acute lymphoblastic leukemia (ALL), where the abnormal sequence is a fusion of two normal genes, i.e., *bcr-abl*. See, e.g., the description of these cancers in International  
20 Patent Publication No. WO94/13793, published June 23, 1994, and incorporated herein by reference for a description of these diseases. In such cancers or diseases, such as CML, the afflicted mammal also possesses a normal copy of the polynucleotide sequence or gene, and the differences between the abnormal and normal sequences or genes are differences in nucleotide sequence. For example, for  
25 CML, the abnormal sequence is the *bcr-abl* fusion, while the normal sequence is *bcr* and *abl*. Thus, the method above can be employed with the target polynucleotide sequence being the sequence which spans the fusion. A method of treatment or prophylaxis of such a cancer in a mammal comprises administering to the mammal a composition of this invention wherein the target polynucleotide is a polynucleotide  
30 sequence of an abnormal cancer-causing gene in a mammal which also possesses a

normal copy of the gene, and wherein the differences between the abnormal gene and the normal gene are differences in polynucleotide sequence. The composition is administered as above, with an optional second agent that facilitates polynucleotide uptake in a cell, and in a pharmaceutically acceptable carrier and in an amount effective to reduce or inhibit the function of the abnormal sequence in the mammal.

The present invention thus encompasses methods for evoking the above-described molecular mechanism for treating any disease or disorder in a mammal characterized by expression of an undesirable polynucleotide product or polynucleotide mediated function not found in a healthy mammal by use of a composition which can deliver to the cells of the mammal the partially double-stranded RNA molecule substantially homologous to the target polynucleotide sequence which expresses or mediates the undesired product or function, in an amount effective to reduce or inhibit the function of that polynucleotide in the cells of the mammal. Provided that the RNA molecule is sufficiently non-homologous to essential mammalian polynucleotide sequences, so that it does not inhibit the function of those essential sequences, this method can be clearly seen to have many therapeutic and prophylactic uses. One of skill in the art can readily select disorders described above, and can also readily select target polynucleotide sequences against which the compositions of the present invention are directed.

#### **D. Other Methods of The Present Invention**

The compositions described above, and the general methods of using these compositions to inhibit or reduce the function of a target polynucleotide sequence, can also be applied to a variety of research, and *in vitro* applications. For example, the method of this invention can be applied to research to determine the function of a selected polynucleotide sequence in a cell line, or a mammalian laboratory animal, by administering to that cell in tissue culture or that animal *in vivo* a composition of the invention wherein the RNA molecule polynucleotide sequence is substantially homologous to the selected sequence and preferably substantially non-homologous to

other polynucleotide sequences in the animal. The inhibition of the function of that target sequence permits study of its influence on the animal's biology and physiology.

Similarly, application of this method can be used to make cell lines of mammalian, bacterial, yeast, fungal, insect and other origins defective in selected pathways by "silencing" a selected functional sequence, such as an enzymatic sequence, a protein expressing sequence, or regulatory sequences necessary to the expression thereof. Such manipulated cells may be employed in conventional assays or drug screening assays, etc.

In an analogous method, a "knock-out" laboratory animal can be prepared by altering the dosage of administration sufficient to permanently shut off the function of a selected gene. Thus, the method of the present invention in delivering an RNA molecule with a polynucleotide sequence sufficiently homologous to the sequence selected to be "knocked out" in the laboratory animal as described above provides a simpler technique for developing "knock-out" mice and other laboratory animals useful for pharmaceutical and genetic research.

Still other research methods for use of the compositions and methods of this invention include the preparation of mutants of microorganisms, both eukaryotic and prokaryotic, for use as research agents or as industrial production strains for the microbial production of desired proteins. Still other uses are expected to be obvious to the person of skill in the art given the teachings herein.

The following examples illustrate methods for preparing the compositions and using the compositions of this invention to reduce or inhibit target polynucleotide sequences. These examples which employ as the agent of the composition, double stranded RNA molecules made by *in vitro* synthesis and target polynucleotide sequences of HIV gag or HSV gD2 merely illustrate embodiments of this invention. It is understood by one of skill in the art, that other selections for the various agents of the compositions, and identity of the target polynucleotide sequences may be readily selected as taught by this specification. These examples are illustrative only and do not limit the scope of the invention.

EXAMPLE 1: REDUCING OR INHIBITING THE FUNCTION OF HIV p24 IN VIRALLY INFECTED CELLS

During the course of HIV infection, the viral genome is reverse transcribed into a DNA template which is integrated into the host chromosome of infected dividing cells. The integrated copy is now a blueprint from which more HIV particles are made. According to this invention, if the function of a polynucleotide sequence essential to replication and/or pathogenesis of HIV is reduced or inhibited, the viral infection can be treated. This example demonstrates the performance of one embodiment of the method of this invention.

10 The plasmid, HIVgpt (AIDS Research and Reference Reagent Program Catalog) was used to generate stable integrated Rhabdomyosarcoma (RD) or COS7 cell lines that contain integrated copies of the defective HIV genome, HIVgpt. The HIVgpt genome encodes a mycophenolic acid (MPA) resistance gene in place of the envelope gene and thereby confers resistance to MPA. The cell lines were made by  
15 transfecting cells with the plasmid followed by selection of cells in MPA. Cells resistant to MPA were clonally amplified. The media from the cultured clonally expanded cells was then assayed for the presence of p24 (an HIV gag polypeptide which is secreted extracellularly) using the p24 ELISA assay kit (Coulter Corporation). All cells were positive for p24.

20 Two RD cell lines and two COS7 cell lines are used to demonstrate one embodiment of the method of the present invention, i.e., reducing or inhibiting the function of the HIV p24 target polynucleotide, which controls p24 synthesis in these cells.

To generate a reagent of the present invention, a 600 polynucleotide (nt) sense  
25 RNA, a 600 nt antisense RNA, and a 600 bp double stranded RNA (dsRNA) mapping to the same coordinates of the gag gene of HIV strain HXB2 and lacking a cap, a poly-adenylation sequence, and a native initiation codon, are used to transfect cells. The RNA molecules are generated through transcription of PCR products bearing a bacteriophage T7 polymerase promoter at one end (see Figs. 1A and 1B). The  
30 coordinates of the primers were derived from the map of the complete genome of

HIV(HXB2), Genbank Accession number K03455 [see also, L. Ratner *et al.*, AIDS Res. Hum. Retroviruses, 3(1):57-69 (1987)]. The Forward gag primer maps to coordinates 901-924 and this sequence follows the T7 promoter in the T7 Forward gag primer. The Reverse gag primer maps to coordinates 1476-1500 and follows the T7 promoter in the T7 Reverse gag primer.

To generate a composition of this invention where the agent is single-stranded sense RNA, a T7 promoter is located at the 5' end of the forward PCR primer. The PCR primers used to generate the DNA template that encodes the ss sense RNA, written 5' to 3' with the top strand of the T7 promoter underlined, are the T7 forward gag primer [SEQ ID NO: 1]:

5' GTAATACGACTCACTATAGGGCGGCAGGGAGCTAGAACGATTCGCAG 3'  
and the Reverse gag primer [SEQ ID NO: 2]:

5' CTGCTATGTCACTTCCCCTTGGTTC 3'

To generate a composition where the agent is a single stranded anti-sense RNA molecule, the T7 promoter is located at the 5' end of the reverse PCR primer. These primers are the T7 Reverse gag primer [SEQ ID NO: 3]:

5' GTAATACGACTCACTATAGGGCGCTGCTATGTCACTTCCCCTTGGTTC 3'  
and the Forward gag primer [SEQ ID NO: 4]:  
5' GCAGGGAGCTAGAACGATTCGCAG 3'.

Both types of PCR products described above are included in the T7 transcription reaction to generate a composition where the agent is double-stranded RNA molecule. Alternatively, an agent of the composition according to this invention is prepared by mixing together sense and anti-sense RNA after transcription.

As a control, similarly sized sense RNA, antisense RNA, and dsRNA molecules are derived from the gD gene of a Herpes Simplex Virus, type 2 genome are generated by the same PCR and T7 transcription techniques. The coordinates of the PCR primers for HSV gD are derived from the map of GenBank Accession number K01408, HSVgD2 gene. The Forward gD primer maps to coordinates 313-336; this sequence follows the T7 promoter in the T7 Forward gD primer. The

Reverse gD primer maps to coordinates 849-872, and follows the T7 promoter in the T7 Reverse gD primer. The primer sets used to generate these control molecules were:

T7 forward gD primer [SEQ ID NO: 5]:

5 5' GTAATACGACTCACTATAGGGCGGTGCGGGTGGGACTCCGCGTCGTC 3'  
and

Forward gD primer [SEQ ID NO: 6]: 5' GTCGCGGTGGGACTCCGCGTCGTC 3';  
and

T7 reverse gD primer [SEQ ID NO: 7]:

10 5' GTAATACGACTCACTATAGGGCGGTGATCTCCGTCCAGTCGTTTATC 3'  
and

Reverse gD primer [SEQ ID NO: 8]: 5' GTGATCTCCGTCCAGTCGTTTATC 3'.

These RNA molecules of the invention and the above-described control molecules are assayed with the RD and COS7 cell lines as follows:

15 Between  $5-6 \times 10^5$  cells/well in six-well plates are cultured to about 80-90% confluence, and are transfected with 2-3  $\mu$ g of a selected RNA molecule or control molecule, using 10  $\mu$ l lipofectamine (Gibco-BRL) as a transfecting agent. Transfected cells are incubated for times ranging between 1 to 17 hours. Another cell culture was transfected with doses of RNA ranging between 1  $\mu$ g to 500  $\mu$ gs, delivered with no  
20 known transfecting agent and incubated on the cells from 0.5 minutes to about two days. For example, one group of cells is transfected with the sense gag RNA, another with the antisense gag RNA, another with ds gag RNA, another with sense gD RNA control, another with antisense gD RNA control, and another with ds gD RNA control. Also additional negative controls are cells which receive no RNA molecules.

25 The cells are cultured at 37°C and monitored for p24 synthesis over the course of several weeks. The cells are assayed three times per week after two days post-administration of RNA, both by measuring p24 in the media of cells using the p24 ELISA assay kit (Coulter Corp) and by immunostaining fixed cells for p24 using a rabbit polyclonal anti-p24 sera (Intracell Corp.) and anti-rabbit IgG that is FITC-  
30 conjugated (Sigma).

According to the present invention, none of the gD RNA molecules demonstrate the ability to retard or inhibit p24 synthesis. However, according to the invention the ds gag RNA inhibits or down regulates p24 synthesis. The sense and antisense RNA molecules are expected to cause only a modest, if any, inhibitory effect on p24 synthesis, unless these RNAs were able to form some degree of double strandedness.

**EXAMPLE 2: DETERMINATION OF THE EXTENT OF REDUCTION OF P24 SYNTHESIS FROM ONE CELL CULTURE TO ANOTHER**

To demonstrate that the down-regulated signal can be transmitted to cells which have not been down-regulated, this example demonstrates that the reduction/inhibition effect (i.e., inhibition or reduction of p24 synthesis) is transmitted to cells in culture that are not transfected by the agent.

*A. Co-Culture of COS 7 and RD cells*

Cells from the cultures of Example 1 which demonstrate reduction of p24 synthesis are co-cultured with control cells of cells that have not previously been incubated with any RNA molecule, and are, in fact, synthesizing p24 at wild-type levels. According to the present invention, the previously transfected cells can transfer the target polynucleotide function inhibition to non-transfected cells, and the control cells in the co-culture are characterized by a reduction in synthesis of p24.

In order to distinguish control cells from the previously transfected cells in the culture, a first protocol is followed: The COS 7 cells of Example 1 which demonstrate inhibition of p24 synthesis are co-cultured with non-transfected RD cells expressing p24 at wildtype levels at various ratios of cell types, e.g., the ratios range from 1/1000 to 1/10 (COS 7/RD) to a total of  $6 - 7 \times 10^5$  cells in 6 well plates. After 2 days of culture under the conditions specified in Example 1, the RD cells in the cultures are examined for p24 synthesis. The cells are examined about 3 times per week for 3 weeks.

p24 synthesis is assayed by two methods. In the first method, the media from the co-cultured cells is assayed for p24 using the p24 ELISA assay



(Coulter). In the second method, cells are immunostained for p24 using rabbit polyclonal sera (Intracell Corp.) against p24 and anti-rabbit IgG conjugated to FITC. Because COS 7 and RD cells are distinguishable by morphology, a loss of stain in the RD cells can be readily distinguished from the COS 7 cells. Because COS 7 cells  
5 express T Antigen while RD cells do not, the co-cultured cells are also stained for T Ag using mouse monoclonal sera against SV40 T antigen (Pharmagen Corp.) and anti-mouse IgG conjugated to r-phycoerythrin (PE). Only the COS 7 cells stain under these conditions. The cell staining is determined by fluorescence microscopy or by FLOW cytometry.

10                   The inhibition of p24 function in RD cells in coculture is demonstrated by comparison to a control culture containing only the RD cells by a loss of FITC stain in the co-cultured RD cells. RD cells in the coculture that are not stained with FITC or PE are evidence of reduction or inhibition of the p24 synthesis function of the p24 target polynucleotide by the RNA molecules (particularly the ds RNA molecules)  
15 of Example 1.

*B. Cultures of transfected RD cells with non-transfected RD cells*

In a second protocol, the transfected RD cells of Example 1, which demonstrate reduced p24 production are co-cultured with non-transfected RD cells which are engineered to contain an integrated hygromycin resistance gene and express  
20 normal levels of p24 using different ratios of cells, with ratios ranging from 1/1000 to 1/10 (RD/control RD) to a total cell number of  $6 - 7 \times 10^5$  in a 6 well plate. Hygromycin-resistant RD cells are made as follows: RD cells ( $5-6 \times 10^5$  cells) are cultured to 80-90% confluence in a six-well plate and are transfected with 2.5  $\mu$ g of the Nru I-Sal I fragment of pCEP4 (Invitrogen Corp.) that contains the hygromycin  
25 resistance gene under the control of a thymidine kinase (TK) promoter. Transfections are done using the transfecting agent, lipofectamine (Gibco BRL). Two days following transfection, the cells are incubated in the presence of 400  $\mu$ g/ml hygromycin. Resistant cells are clonally expanded. One or more of the clonally expanded cell lines are used as the control in the experiment.

From 1 day to several weeks after co-culture under the conditions specified in Example 1, replicate co-cultures are incubated with 400 µg/ml hygromycin. This concentration of hygromycin kills the RD cells that are not hygromycin resistant, leaving only the control hygromycin resistant RD cells. The remaining resistant cells are derived from the control cells. P24 levels are measured directly from the control cells, for example using the ELISA of Example 1 as well as by immunostaining as above described.

According to the present invention, inhibition of p24 production is revealed in at least a subset of the control cells.

10 EXAMPLE 3: *IN VIVO* INHIBITION OF ENDOGENOUS INTERLEUKIN-12 PRODUCTION BY THE METHOD OF THIS INVENTION

A. *Design of RNA molecules as Compositions of the Invention*

All RNA molecules in this experiment are close to 600 nts in length, and all RNA molecules are designed to be incapable of producing the p40 chain of IL-12. The molecules have no cap and no poly-A sequence; the native initiation codon is not present, and the RNA does not encode the full-length product. The following RNA molecules are designed:

- (1) a single-stranded (ss) sense RNA polynucleotide sequence homologous to IL-12 p40 murine messenger RNA (mRNA);
- 20 (2) a ss anti-sense RNA polynucleotide sequence complementary to IL-12 p40 murine mRNA,
- (3) a double-stranded (ds) RNA molecule comprised of both sense and anti-sense p40 IL-12 murine mRNA polynucleotide sequences,
- (4) a ss sense RNA polynucleotide sequence homologous to IL-12 p40 murine heterogeneous RNA (hnRNA),
- 25 (5) a ss anti-sense RNA polynucleotide sequence complementary to IL-12 p40 murine hnRNA,
- (6) a ds RNA molecule comprised of the sense and anti-sense IL-12 p40 murine hnRNA polynucleotide sequences,

(7) a ss murine RNA polynucleotide sequence homologous to the top strand of the IL-12 p40 promoter,

(8) a ss murine RNA polynucleotide sequence homologous to the bottom strand of the IL-12 p40 promoter, and

5 (9) a ds RNA molecule comprised of murine RNA polynucleotide sequences homologous to the top and bottom strands of the IL-12 p40 promoter.

As a negative control the sense, anti-sense and ds RNAs derived from the HSV2 gD gene described in Example 1 are also used. Another control group is composed of mice receiving no RNA.

10 As described in Example 1, the various RNA molecules of (1)-(9) above are generated through T7 RNA polymerase transcription of PCR products bearing a T7 promoter at one end. In the instance where a sense RNA is desired, a T7 promoter is located at the 5' end of the forward PCR primer. In the instance where an antisense RNA is desired, the T7 promoter is located at the 5' end of the reverse  
15 PCR primer. When dsRNA is desired both types of PCR products are included in the T7 transcription reaction. Alternatively, sense and anti-sense RNA are mixed together after transcription.

The PCR primers used in the construction of the RNA molecules of this Example are 5' to 3', with the top strand of the T7 promoter underlined.

20 Forward IL-12 genomic (hnRNA) [SEQ ID NO: 9]:  
5' TCAGCAAGCACTTGCCAAACTCCTG 3' and Reverse IL-12 genomic (hnRNA) [SEQ ID NO: 10]: 5' GAGACAAGGTCTCTGGATGTTATTG 3';  
T7 Forward IL-12 genomic (hnRNA) [SEQ IDNO: 11]:  
5' GTAATACGACTCACTATAGGGTCAGCAAGCACTTGCCAAACTCCTG 3'  
25 and T7 Reverse IL-12 genomic (hnRNA) [SEQ ID NO: 12]:  
5' GTAATACGACTCACTATAGGGGAGACAAGGTCTCTGGATGTTATTG 3';  
T7 Forward IL-12 promoter [SEQ ID NO: 13]:  
5' GTAATACGACTCACTATAGGGCCTATAAGCATAAGAGACGCCCTC 3' and  
Forward IL-12 promoter [SEQ ID NO: 14]:  
30 5' CCTATAAGCATAAGAGACGCCCTC 3';

Reverse IL-12 promoter [SEQ ID NO: 15]:

5' GGCTGCTCCTGGTGCTTATATAC 3'

and T7 Reverse IL-12 promoter [SEQ ID NO: 16]:

5' GTAATACGACTCACTATAGGGGGCTGCTCCTGGTGCTTATATAC 3';

5 T7 Forward IL-12 cDNA (mRNA) [SEQ ID NO: 17]:

5' GTAATACGACTCACTATAGGGTGTGTCCTCAGAAGCTAACCATC 3' and

Forward IL-12 cDNA (mRNA) [SEQ ID NO: 18]:

5' TGTGTCCTCAGAAGCTAACCATC 3';

Reverse IL-12 cDNA (mRNA) [SEQ ID NO: 19]:

10 5' GCAGGTGACATCCTCCTGGCAGGA 3'

and T7 Reverse IL-12 cDNA (mRNA) [SEQ ID NO: 20]:

5' GTAATACGACTCACTATAGGGGCAGGTGACATCCTCCTGGCAGGA 3'.

The genomic and PCR primer coordinates are based on the map supplied in the following citation: Tone *et al*, Eur. J. Immunol., 26:1222-1227

15 (1996). The forward IL-12 genomic primer maps to coordinates 8301-8325. The reverse IL-12 genomic primer maps to coordinates 8889-8913. The forward IL-12 promoter primer maps to coordinates 83-106. The reverse IL-12 promoter primer maps to coordinates 659-682. The coordinates for the cDNA PCR primers is based on GenBank Accession No. M86671. The forward IL-12 cDNA primer maps to  
20 coordinates 36-58. The reverse IL-12 cDNA primer maps to coordinates 659-682.

#### B. Assay

Balb/c mice (5 mice/group) are injected intramuscularly or intraperitoneally with the murine IL-12 p40 chain specific RNAs described above or with controls identified above at doses ranging between 10 µg and 500 µg. Sera is  
25 collected from the mice every four days for a period of three weeks and assayed for IL-12 p40 chain levels using the Quantikine M-IL-12 p40 ELISA Assay (Genzyme).

According to the present invention, mice receiving ds RNA molecules derived from both the IL-12 mRNA, IL-12 hnRNA and ds RNA derived from the IL-12 promoter demonstrate a reduction or inhibition in IL-12 production. A modest, if

any, inhibitory effect is observed in sera of mice receiving the single stranded IL-12 derived RNA molecules, unless the RNA molecules have the capability of forming some level of double-strandedness. None of the HSV gD derived RNAs are expected to reduce or inhibit IL-12 *in vivo* in a specific manner..

5    EXAMPLE 4: METHOD OF THE INVENTION IN THE PROPHYLAXIS OF DISEASE

          A.    *In Vitro Assay*

          Vero and/or BHK cells, seeded at a density of 20-30% confluency, are cultured in six-well plates at 37°C in DMEM with 10% FBS. When cells are 80-90%  
10    confluent, they are transfected with 2-3 µg of the HIV gag- and HSV gD- specific RNA molecules described in Example 1 using lipofectamine (Gibco-BRL) as a transfecting agent. The RNA molecules are also delivered in the absence of any known transfecting agent in amounts varying between 5 and 100 µg. Another group of cells receives no RNA.

15                Still other groups of Vero and/or BHK cells are similarly transfected with 2-3 µg of a double-stranded DNA plasmid, plasmid 24, which is described in U. S. Patent No. 5,851,804, incorporated herein by reference, which contains a sequence encoding the HSV2 gD protein under the control of the HCMV promoter and a SV40 polyA sequence.

20                The transfected cells are cultured at 37°C in DMEM with 10% FBS. At days 1, 2, 4 and 7 following transfection, cells are infected with HSV2 at a multiplicity of infection (MOI) of 0.1 in an inoculum of 250 µl DMEM. The inoculum is allowed to adsorb for 1 hour after which 2 mls of DMEM (10% FBS) is added per well. For those cells infected at 4 and 7 days post transfection, the cells are passaged  
25    into a new six-well plate such that they are confluent at the time of infection. If the cells are not passaged, they become overcrowded.

          At 36-48 hours post-infection, the cell lysates are assayed for viral titer by conventional plaque assay on Vero cells [Clinical Virology Manual, 2d edit., eds. S. Specter and G. Lancz, pp. 473-94 (1992)]. According to this invention, the cells

transfected with the ds DNA plasmid, APL-400-024, and with the ds RNA molecule containing a polynucleotide sequence of the gD2 antigen, cannot be productively infected with HSV2. All other cells are anticipated to become productively infected with HSV2.

5           B.     *In Vivo Assay*

Using the HSV-gD specific RNA molecules described in Example 1, which do not have the ability to make HSVgD protein and HIV gag specific RNA molecules as controls, mice are evaluated for protection from HSV challenge through the use of the injected HSVgD specific RNA molecules of the invention.

10           Balb/c mice (5 mice/group) are immunized intramuscularly or intraperitoneally with the described RNA molecules at doses ranging between 10 and 500 µg RNA. At days 1, 2, 4 and 7 following RNA injection, the mice are challenged with HSV-2 ( $10^5$  pfu in 30 µls) by intravaginal inoculation. Everyday post HSV-2 inoculation, the mice are observed for signs of infection and graded on a scale of 0-4.  
15     Zero is no sign of infection; 1 denotes redness; 2 denotes vesicles and redness; 3 denotes vesicles, redness and incontinence; and 4 denotes paralysis.

According to the present invention, because the mice that receive dsRNA molecules of the present invention which contain the HSV gD sequence are shown to be protected against challenge. The mice receiving the HIV gag control  
20     RNA molecules are not protected. Mice receiving the ss RNA molecules which contain the HSV gD sequence are expected to be minimally, if at all, protected, unless these molecules have the ability to become at least partially double stranded *in vivo*. According to this invention, because the dsRNA molecules of the invention do not have the ability to make HSV gD protein, the protection provided by delivery of the  
25     RNA molecules to the animal is due to a non-immune mediated mechanism that is gene specific.

All above-noted published references are incorporated herein by reference.  
Numerous modifications and variations of the present invention are included in the  
above-identified specification and are expected to be obvious to one of skill in the art.  
Such modifications and alterations to the compositions and processes of the present  
5 invention are believed to be encompassed in the scope of the claims appended hereto.

## WHAT IS CLAIMED IS:

1. A composition for inhibiting the function of a target polynucleotide sequence in a mammalian cell, wherein said composition comprises an agent that provides to a mammalian cell an at least partially double-stranded RNA molecule that does not produce a functional protein, and that comprises a polynucleotide sequence of at least about 200 nucleotides in length, said polynucleotide sequence being substantially homologous to said target polynucleotide sequence, and substantially non-homologous to a selected naturally-occurring, essential mammalian polynucleotide sequence.
2. The composition according to claim 1 wherein at least 11 contiguous nucleotides of said polynucleotide sequence of said RNA molecule are present in a double-stranded sequence, depending upon the composition of said polynucleotide sequence and a  $\Delta G$  of about -9.2 kcal/mol.
3. The composition according to claim 2 wherein substantially the entire polynucleotide sequence of said RNA molecule is double stranded.
4. The composition according to claim 1 wherein said RNA molecule polynucleotide sequence has a sequence of between at least about 12 to about 16 contiguous nucleotides in exact homology to said target polynucleotide sequence, and wherein said overall homology of said RNA molecule polynucleotide sequence to said target sequence is greater than about 10%.
5. The composition according to claim 4, wherein said homology is greater than about 50%.



6. The composition according to claim 1 wherein said agent is an RNA molecule made by enzymatic synthetic methods or chemical synthetic methods *in vitro*.

7. The composition according to claim 1 wherein said agent is an RNA molecule made *in vitro* by isolation from a recombinant microorganism or the culture media in which said microorganism is grown.

8. The composition according to claim 1 wherein said agent generates said RNA molecule *in vivo* after delivery to said mammalian cell.

9. The composition according to claim 1 wherein said agent is a double stranded RNA.

10. The composition according to claim 1 wherein said agent is a single stranded RNA sense strand.

11. The composition according to claim 10 wherein said single stranded RNA sense strand forms a hairpin at one or both termini or intermediate between said termini.

12. The composition according to claim 10 wherein said single stranded RNA sense strand folds back upon itself to become partially double stranded.

13. The composition according to claim 1 wherein said agent is a single stranded RNA anti-sense strand.

14. The composition according to claim 13 wherein said single stranded RNA anti-sense strand forms a hairpin at one or both termini or intermediate between said termini.

15. The composition according to claim 13 wherein said single stranded RNA anti-sense strand folds back upon itself to become partially double stranded.
16. The composition according to claim 1, wherein said agent is a single stranded RNA sequence comprising both a sense polynucleotide sequence and an anti-sense polynucleotide sequence, optionally separated by a non-base paired polynucleotide sequence, said single stranded RNA sequence having the ability to become double-stranded.
17. The composition according to claim 1 wherein said agent is a circular RNA molecule that forms a rod structure.
18. The composition according to claim 8 wherein said agent is a double stranded DNA molecule encoding said RNA molecule.
19. The composition according to claim 18 wherein said DNA encodes a double stranded RNA.
20. The composition according to claim 18 wherein said DNA encodes a single stranded RNA sense strand.
21. The composition according to claim 20 wherein said DNA encodes a single stranded RNA sense strand that forms a hairpin at one or both termini or intermediate therebetween.
22. The composition according to claim 20 wherein said DNA encodes a single stranded RNA sense strand that folds back upon itself to become partially double stranded.

23. The composition according to claim 18 wherein said DNA encodes a single stranded RNA anti-sense strand.

24. The composition according to claim 23 wherein said DNA encodes a single stranded RNA anti-sense strand that forms a hairpin at one or both termini or intermediate therebetween.

25. The composition according to claim 23 wherein said DNA encodes a single stranded RNA anti-sense strand that folds back upon itself to become partially double stranded.

26. The composition according to claim 18 wherein said DNA encodes a single stranded RNA sequence comprising both a sense polynucleotide sequence and an anti-sense polynucleotide sequence, optionally separated by a non-base paired polynucleotide sequence, said single stranded RNA sequence having the ability to become double-stranded.

27. The composition according to claim 18 wherein said DNA encodes a circular RNA molecule that forms a rod structure.

28. The composition according to claim 1, wherein said agent is a plasmid.

29. The composition according to claim 1, wherein said agent comprises a first DNA plasmid encoding a single stranded RNA sense polynucleotide sequence and a second DNA plasmid encoding a single stranded RNA anti-sense polynucleotide sequence, wherein said sense and anti-sense RNA sequences have the ability to base-pair and become double-stranded.

30. The composition according to claim 28, wherein said plasmid comprises bacterial sequences.

31. The composition according to claim 1, wherein said agent is a recombinant bacterium.

32. The composition according to claim 1, wherein said agent is a recombinant virus.

33. The composition according to claim 1, wherein said agent is a donor cell transfected *in vitro* with the molecule described in any of claims 2 through 32.

34. The composition according to any of claims 30-32, wherein said agent is selected from the group consisting of a living recombinant virus or bacteria or cell, a dead virus or bacteria or cell, or an inactivated virus or bacteria or cell.

35. The composition according to claim 1, wherein said agent lacks a polyadenylation sequence.

36. The composition according to claim 1, wherein said RNA molecule is not translated.

37. The composition according to claim 1, wherein said agent lacks a Kozak region.

38. The composition according to claim 1, wherein said agent lacks an initiating methionine codon.

39. The composition according to claim 1 wherein said RNA molecule lacks a cap structure.

40. The composition according to claim 1 wherein said agent lacks signals for protein synthesis.

41. The composition according to claim 1, comprising a mixture of different said agents.

42. The composition according to claim 1 wherein said target polynucleotide sequence is a virus polynucleotide sequence necessary for replication and/or pathogenesis of said virus in an infected mammalian cell.

43. The composition according to claim 42, wherein said virus is selected from the group consisting of a DNA virus and a virus that has an intermediary DNA stage.

44. The composition according to claim 43, wherein said virus is selected from the group consisting of Retrovirus, Herpesvirus, Hepadenovirus, Poxvirus, Parvovirus, Papillomavirus, and Papovavirus.

45. The composition according to claim 44, wherein said virus is selected from the group consisting of HIV, HBV, HSV, CMV, HPV, HTLV and EBV.

46. The composition according to claim 1, wherein said target polynucleotide sequence is a tumor antigen or functional fragment thereof or a regulatory sequence of a virus-induced cancer, which antigen or sequence is required for the maintenance of said tumor in said mammal.

47. The composition according to claim 46, wherein said cancer is selected from the group consisting of HPV E6/E7 virus-induced cervical carcinoma, HTLV-induced cancer and EBV induced cancer.

48. The composition according to claim 1, wherein said target polynucleotide sequence is a polynucleotide sequence of an intracellular or extracellular pathogen necessary for replication and/or pathogenesis of said pathogen in an infected mammalian cell.

49. The composition according to claim 1 wherein said target polynucleotide sequence is a polynucleotide sequence of an abnormal cancer-causing sequence in a mammal which also possesses a normal copy of said sequence, and wherein the differences between the abnormal and the normal sequences are differences in polynucleotides.

50. The composition according to claim 49 wherein said abnormal sequence is a fusion of two normal genes.

51. The composition according to claim 50 wherein said target polynucleotide is the polynucleotide sequence spanning said fusion.

52. A pharmaceutical composition comprising a composition of any of claims 1-51, and an optional second agent that facilitates polynucleotide uptake in a cell, in a pharmaceutically acceptable carrier.

53. The composition according to claim 52, wherein said second agent is selected from the group consisting of a local anaesthetic, a peptide, a lipid including cationic lipids, a liposome or lipidic particle, a polycation, a branched, three-dimensional polycation, a carbohydrate, a cationic amphiphile, a detergent, a benzylammonium surfactant, or another compound that facilitates polynucleotide transfer to cells.

54. The composition according to claim 53 wherein said second agent is bupivacaine.

55. A method for treating a viral infection in a mammal, comprising:  
administering to said mammal a composition according to claim 1, with  
an optional second agent that facilitates polynucleotide uptake in a cell, in a  
pharmaceutically acceptable carrier, wherein said target polynucleotide is a virus  
polynucleotide sequence necessary for replication and/or pathogenesis of said virus in  
an infected mammalian cell, in an amount effective to reduce or inhibit the function of  
said viral sequence in the cells of said mammal.

56. A method for preventing a viral infection in a mammal, comprising:  
administering to said mammal a composition according to claim 1, with  
an optional second agent that facilitates polynucleotide uptake in a cell, in a  
pharmaceutically acceptable carrier, wherein said target polynucleotide is a virus  
polynucleotide sequence necessary for replication and/or pathogenesis of said virus in  
an infected mammalian cell, in an amount effective to reduce or inhibit the function of  
said viral sequence upon subsequent introduction of said virus into said mammalian  
cells.

57. A method for treatment or prophylaxis of a virally induced cancer in a  
mammal comprising:  
administering to said mammal a composition according to claim 1, with  
an optional second agent that facilitates polynucleotide uptake in a cell, in a  
pharmaceutically acceptable carrier, wherein said target polynucleotide is a sequence  
encoding a tumor antigen, a regulatory sequence, or a functional fragment thereof,  
which antigen or sequence function is required for the maintenance of said tumor in  
said mammal, in an amount effective to reduce or inhibit the function of said antigen in  
said mammal.

58. A method for the treatment or prophylaxis of infection of a mammal by an intracellular or extracellular pathogen comprising administering to said mammal a composition according to claim 1, with an optional second agent that facilitates polynucleotide uptake in a pathogenic or mammalian cell, in a pharmaceutically acceptable carrier, wherein said target polynucleotide is a polynucleotide sequence of said pathogen necessary for replication and/or pathogenesis of said pathogen in an infected mammal or mammalian cell, in an amount effective to reduce or inhibit the function of said sequence in said mammal.

59. A method of treatment or prophylaxis of cancer in a mammal comprising administering to said mammal a composition according to claim 1, with an optional second agent that facilitates polynucleotide uptake in a cell, in a pharmaceutically acceptable carrier, wherein said target polynucleotide is a polynucleotide sequence of an abnormal cancer-causing sequence in a mammal which also possesses a normal copy of said sequence, and wherein the differences between the abnormal sequence and said normal sequence are differences in polynucleotides, in an amount effective to reduce or inhibit the function of said abnormal sequence in said mammal.

60. A method for treating a disease or disorder in a mammal comprising:  
administering to said mammal having a disease or disorder characterized by expression of polynucleotide product not found in a healthy mammal, a composition according to claim 1, wherein said target polynucleotide sequence is a polynucleotide sequence which expresses said polynucleotide product or regulatory sequence necessary to expression of said product, in an amount effective to reduce or inhibit the function of said target polynucleotide product in the cells of said mammal.



61. Use of a composition according to claim 1, with an optional second agent that facilitates polynucleotide uptake in a cell, in a pharmaceutically acceptable carrier, wherein said target polynucleotide is a virus polynucleotide sequence necessary for replication and/or pathogenesis of said virus in an infected mammalian cell, in the preparation of a medicament for treating a viral infection in a mammal.

62. Use according to claim 61, wherein said composition is in an amount effective to reduce or inhibit the function of said viral sequence in the cells of said mammal.

63. Use according to claim 61, wherein said composition is in an amount effective to reduce or inhibit the function of said viral sequence upon subsequent introduction of said virus into said mammalian cells.

64. Use of a composition according to claim 1, with an optional second agent that facilitates polynucleotide uptake in a cell, in a pharmaceutically acceptable carrier, wherein said target polynucleotide is a sequence encoding a tumor antigen, a regulatory sequence, or a functional fragment thereof, which antigen or sequence function is required for the maintenance of said tumor in said mammal, in an amount effective to reduce or inhibit the function of said antigen in said mammal, in the preparation of a medicament for treatment or prophylaxis of a virally induced cancer in a mammal.

65. Use of a composition according to claim 1, with an optional second agent that facilitates polynucleotide uptake in a pathogenic or mammalian cell, in a pharmaceutically acceptable carrier, wherein said target polynucleotide is a polynucleotide sequence of said pathogen necessary for replication and/or pathogenesis of said pathogen in an infected mammal or mammalian cell, in an amount effective to reduce or inhibit the function of said sequence in said mammal, in the preparation of a medicament for the treatment or prophylaxis of infection of a mammal by an intracellular or extracellular pathogen.

66. Use of a composition according to claim 1, with an optional second agent that facilitates polynucleotide uptake in a cell, in a pharmaceutically acceptable carrier, wherein said target polynucleotide is a polynucleotide sequence of an abnormal cancer-causing sequence in a mammal which also possesses a normal copy of said sequence, and wherein the differences between the abnormal sequence and said normal sequence are differences in polynucleotides, in an amount effective to reduce or inhibit the function of said abnormal sequence in said mammal, in the preparation of a medicament for the treatment or prophylaxis of cancer in a mammal.

67. Use of a composition according to claim 1, wherein said target polynucleotide sequence is a polynucleotide sequence which expresses said polynucleotide product or regulatory sequence necessary to expression of a polynucleotide product not found in a healthy mammal, in an amount effective to reduce or inhibit the function of said target polynucleotide product in the cells of said mammal, in the preparation of a medicament for treating a disease or disorder in a mammal characterized by expression of said product.

Fig. 1A





Fig. 1B



## SEQUENCE LISTING

<110> American Home Products Corporation  
Pachuk, Catherine  
Satishchandran, C.

<120> Methods and Compositions for Inhibiting the Function of  
Polynucleotide Sequences

<130> AHP28APCT

<140>

<141>

<150> 60/130,377

<151> 1999-04-21

<160> 20

<170> PatentIn Ver. 2.1

<210> 1

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T7 forward  
gag primer

<400> 1

gtaatacgac tcactatagg gcggcaggga gctagaacga ttcgcag 47

<210> 2

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: reverse gag  
primer

<400> 2

ctgctatgtc acttccccctt gggtc 25

<210> 3

<211> 48

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T7 reverse  
gag primer

<400> 3

gtaatacgac tcactatagg gcgctgctat gtcacttccc cttgggtc

48

<210> 4

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: forward gag  
primer

<400> 4

gcagggagct agaacgattc gcag

24

<210> 5

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T7 forward gD  
primer

<400> 5

gtaatacgac tcactatagg gcggtcgagg tgggactccg cgtcgtc

47

<210> 6

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: forward gD  
primer

<400> 6

gtcgcggtgg gactccgct cgtc

24

<210> 7

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T7 reverse gD  
primer

<400> 7

gtaatacgac tcactatagg gcggtgatct ccgtccagtc gtttatac

47

<210> 8

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: reverse gD  
primer

<400> 8

gtgatctccg tccagtcggt tacc

24

<210> 9

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: forward IL-12  
genomic

<400> 9

tcagcaagca cttgccaaac tccg

25

<210> 10

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: reverse IL-12  
genomic

<400> 10

gagacaaggt ctctggatgt tattg

25

<210> 11

<211> 46

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T7 forward  
IL-12 genomic

<400> 11

gtaatacgac tcactatagg gtcagcaagc acttgccaaa ctcttg

46

<210> 12

<211> 46

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T7 reverse  
IL-12 genomic

<400> 12

gtaatacgac tcactatagg ggagacaagg tctctggatg ttattg

46

<210> 13

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T7 forward  
IL-12 primer

<400> 13

gtaatacgac tcactatagg gcctataagc ataagagacg ccctc

45

<210> 14

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: forward IL-12 promoter

<400> 14

cctataagca taagagacgc cctc

24

<210> 15

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: reverse IL-12 promoter

<400> 15

ggctgctcct ggtgcttata tac

23

<210> 16

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T7 reverse IL-12 promoter

<400> 16

gtaatacgac tcactatagg gggctgctcc tgggtgcttat atac

44

<210> 17

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: T7 forward IL-12 cDNA

<400> 17

gtaatacgac tcactatagg gtgtgtcctc agaagctaac catc

44



<210> 18  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: forward IL-12  
cDNA

<400> 18  
tgtgtcctca gaagctaacc atc 23

<210> 19  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: reverse IL-12  
cDNA

<400> 19  
gcaggtgaca tcctcctggc agga 24

<210> 20  
<211> 45  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: T7 reverse  
IL-12 cDNA

<400> 20  
gtaatacgac tcactatagg ggcaggtgac atcctcctgg cagga 45

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
26 October 2000 (26.10.2000)

PCT

(10) International Publication Number  
WO 00/63364 A3

(51) International Patent Classification?: C12N 15/11,  
15/63, 15/70, 15/86, C07K 14/01, 14/08, A61K 48/00,  
38/16, A61P 35/00

(21) International Application Number: PCT/US00/10555

(22) International Filing Date: 19 April 2000 (19.04.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/130,377 21 April 1999 (21.04.1999) US

(71) Applicant (for all designated States except US): AMER-  
ICAN HOME PRODUCTS CORPORATION [US/US];  
Five Giralda Farms, Madison, NJ 07940 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PACHUK, Cath-  
erine [US/US]; 3212 New Park Circle, Lansdale, PA 19446  
(US). SATISHCHANDRAN, C. [US/US]; 605 Shepard  
Drive, Lansdale, PA 19446 (US).

(74) Agents: BAK, Mary, E. et al.; Howson and Howson,  
Spring House Corporate Center, P.O. Box 457, Spring  
House, PA 19477 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,  
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,  
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,  
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ,  
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,  
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- With international search report.
- Before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments.

(88) Date of publication of the international search report:  
11 January 2001

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: METHODS AND COMPOSITIONS FOR INHIBITING THE FUNCTION OF POLYNUCLEOTIDE SEQUENCES

(57) Abstract: A therapeutic composition for inhibiting the function of a target polynucleotide sequence in a mammalian cell includes an agent that provides to a mammalian cell an at least partially double-stranded RNA molecule comprising a polynucleotide sequence of at least about 200 nucleotides in length, said polynucleotide sequence being substantially homologous to a target polynucleotide sequence. This RNA molecule desirably does not produce a functional protein. The agents useful in the composition can be RNA molecules made by enzymatic synthetic methods or chemical synthetic methods *in vitro*; or made in recombinant cultures of microorganisms and isolated therefrom, or alternatively, can be capable of generating the desired RNA molecule *in vivo* after delivery to the mammalian cell. In methods of treatment of prophylaxis of virus infections, other pathogenic infections or certain cancers, these compositions are administered in amounts effective to reduce or inhibit the function of the target polynucleotide sequence, which can be of pathogenic origin or produced in response to a tumor or other cancer, among other sources.

WO 00/63364 A3

# INTERNATIONAL SEARCH REPORT

International Application No. <b>PCT/US 00/10555</b>				
<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 C12N15/11 C12N15/63 C12N15/70 C12N15/86 C07K14/01 C07K14/08 A61K48/00 A61K38/16 A61P35/00				
According to International Patent Classification (IPC) or to both national classification and IPC				
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N A61K				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, BIOSIS				
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>				
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X	MONTGOMERY M K ET AL: "Double-stranded RNA as a mediator in sequence-specific genetic silencing and co-suppression" TRENDS IN GENETICS, NL, ELSEVIER SCIENCE PUBLISHERS B.V. AMSTERDAM, vol. 14, no. 7, 1 July 1998 (1998-07-01), pages 255-258, XP004124680 ISSN: 0168-9525 the whole document	1		
X	WO 95 27783 A (JOSHI SUKHWAL SADNA) 19 October 1995 (1995-10-19)  page 8, line 29 -page 11, line 10 <div style="text-align: center;">--- -/-</div>	1,4,5,8, 9,28, 32-40, 42-45, 48,55, 56,58, 61-63		
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</span> <span><input checked="" type="checkbox"/> Patent family members are listed in annex.</span> </div>				
* Special categories of cited documents : <div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*8* document member of the same patent family</p> </div> </div>				
Date of the actual completion of the international search <div style="text-align: center;">30 October 2000</div>		Date of mailing of the international search report <div style="text-align: center;">15/11/2000</div>		
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center;">Mateo Roseil, A.M.</div>		

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/10555

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 98 05770 A (ROTHBARTH KARSTEN ;JOSWIG GABY (DE); WERNER DIETER (DE); SCHUBERT) 12 February 1998 (1998-02-12) cited in the application</p> <p>the whole document</p>	<p>1,6,8, 18,20, 23-25, 28,30, 31, 33-40, 49,59, 60,64</p>
X	<p>WO 97 34638 A (UNIV CALIFORNIA) 25 September 1997 (1997-09-25)</p> <p>page 3, line 5 -page 4, line 28; examples 1,2</p>	<p>1,6,8, 13,28, 32-34, 42-45, 48,55-67</p>
X	<p>PLAYER MARK R ET AL: "Targeting HIV mRNA for degradation: 2,5-A antisense chimeras as potential chemotherapeutic agents for AIDS." NUCLEOSIDES &amp; NUCLEOTIDES, vol. 16, no. 7-9, July 1997 (1997-07), pages 1221-1222, XP000952972 ISSN: 0732-8311 the whole document</p>	<p>1,42-45</p>
A	<p>WO 97 07825 A (UNIV COLORADO) 6 March 1997 (1997-03-06)</p> <p>page 3, line 5 -page 5, line 20 page 8, line 3-23</p>	<p>1,6,8, 10,17, 28,33, 34,45</p>
A	<p>US 4 891 315 A (WEIS JOHN H ET AL) 2 January 1990 (1990-01-02) the whole document</p>	<p>1,42,45</p>
A	<p>MIKITA T ET AL: "EFFECTS OF ARABINOSYLCYTOSINE-SUBSTITUTED DNA ON DNA/RNA HYBRID STABILITY AND TRANSCRIPTION BY T7 RNA POLYMERASE" BIOCHEMISTRY,US,AMERICAN CHEMICAL SOCIETY. EASTON, PA, vol. 33, 1994, pages 9195-9208, XP000882431 ISSN: 0006-2960 the whole document</p>	<p>1</p>

-/--

## INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/US 00/10555

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	YUKIKO TONE ET AL: "STRUCTURE AND CHROMOSOMAL LOCATION OF THE MOUSE INTERLEUKIN-12 P35 AND P40 SUBUNIT GENES" EUROPEAN JOURNAL OF IMMUNOLOGY, DE, WEINHEIM, vol. 26, no. 6, 1 June 1996 (1996-06-01), pages 1222-1227, XP000612644 ISSN: 0014-2980 cited in the application the whole document	1
A	--- JAEGER J A ET AL: "IMPROVED PREDICTIONS OF SECONDARY STRUCTURES FOR RNA" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES, vol. 86, no. 20, 1989, pages 7706-7710, XP000953154 1989 ISSN: 0027-8424 cited in the application the whole document	2
A	--- WATERHOUSE ET AL: "Virus resistance and gene silencing in plants can be induced by simultaneous expression of sense and antisense RNA" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE, WASHINGTON, US, vol. 95, November 1998 (1998-11), pages 13959-13964, XP002114472 ISSN: 0027-8424 the whole document	1,42
A	--- NELLEN W ET AL: "WHAT MAKES AN MRNA ANTI-SENSE-ITIVE?" TIBS TRENDS IN BIOCHEMICAL SCIENCES, EN, ELSEVIER PUBLICATION, CAMBRIDGE, vol. 18, no. 11, 1 November 1993 (1993-11-01), pages 419-423, XP002049385 ISSN: 0968-0004 the whole document	1
A	--- FIRE A ET AL: "Potent and specific genetic interference by double-stranded RNA in Caenorhabditis elegans" NATURE, GB, MACMILLAN JOURNALS LTD. LONDON, vol. 391, 19 February 1998 (1998-02-19), pages 806-811, XP002095876 ISSN: 0028-0836 the whole document	1
	---	

-/--

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/10555

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>WIANNY F ET AL: "Specific interference with gene function by double-stranded RNA in early mouse development" NATURE CELL BIOLOGY,GB,MACMILLAN PUBLISHERS, vol. 2, no. 2, February 2000 (2000-02), pages 70-75, XP002138445 ISSN: 1465-7392 the whole document</p> <p>---</p>	1,3,6,8, 9
P,X	<p>WO 99 53050 A (WANG MING BO ;COMMW SCIENT IND RES ORG (AU); GRAHAM MICHAEL WAYNE) 21 October 1999 (1999-10-21) cited in the application</p> <p>page 1, line 1-17 page 8, line 5 -page 10, line 20 page 22, line 4 -page 23, line 2 page 26, line 1-25</p> <p>---</p>	1,3-6,8, 10-16, 35-40, 42, 61-63,65
E	<p>WO 00 44914 A (FARRELL MICHAEL J ;LI YIN XIONG (US); KIRBY MARGARET L (US); MEDIC) 3 August 2000 (2000-08-03) page 2, line 20 -page 5, line 10 page 10, line 4 -page 19, line 17; example III</p> <p>---</p>	1,3,6,8, 9,28,33, 34,42
T	<p>CAPLEN N J ET AL: "dsRNA-mediated gene silencing in cultured Drosophila cells: a tissue culture model for the analysis of RNA interference" GENE,ELSEVIER BIOMEDICAL PRESS. AMSTERDAM,NL, vol. 252, no. 1-2, 11 July 2000 (2000-07-11), pages 95-105, XP004210158 ISSN: 0378-1119 abstract page 96, left-hand column, paragraph 2 page 102, left-hand column, last paragraph -page 104, left-hand column, last paragraph</p> <p>---</p>	1,9,28, 31,33,34
T	<p>PARK WEE-SUNG ET AL: "Inhibition of HIV-1 replication by a new type of circular dumbbell RNA/DNA chimeric oligonucleotides." BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 270, no. 3, 21 April 2000 (2000-04-21), pages 953-960, XP002151277 ISSN: 0006-291X the whole document</p> <p>-----</p>	1,6,8, 42-45, 55,56, 58, 61-63,65

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1,42,46,48,49,52-63

Present claims 1,42,46,48,49 and 52-63 relate to an extremely large number of possible compounds (agents and/or target polynucleotides). In fact, the claims contain so many options that a lack of clarity and conciseness within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible. Consequently, the search has been carried out for those parts of the application which do appear to be clear and concise, namely, (a) agents: partially ds-RNA, dsRNA, ss-RNA sense and antisense, circular RNA, lariats and hybrid RNA-DNA as mentioned in the description at pages 8, line 5 to page 9, line 4, and page 14, line 1 to page 16, line 22, and (b) target polynucleotides gag gene of HIV, the gD gene of HSV, type 2 genome and to a polynucleotide sequence homologous to IL-12p40 murine messenger RNA as recited in examples 1-4.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/10555

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9527783 A	19-10-1995	AU 2133695 A	30-10-1995
WO 9805770 A	12-02-1998	DE 19631919 A	12-02-1998
		EP 0918853 A	02-06-1999
WO 9734638 A	25-09-1997	AU 2585197 A	10-10-1997
WO 9707825 A	06-03-1997	US 5766903 A	16-06-1998
		AU 7103796 A	19-03-1997
		CA 2229454 A	06-03-1997
		EP 0846002 A	10-06-1998
US 4891315 A	02-01-1990	AT 59679 T	15-01-1991
		CA 1295561 A	11-02-1992
		DE 3382072 D	07-02-1991
		DK 333183 A	21-01-1984
		EP 0101655 A	29-02-1984
		ES 524245 D	01-04-1985
		ES 8504254 A	01-07-1985
		ES 529225 D	16-05-1985
		ES 8505253 A	01-09-1985
		FI 832632 A,B,	21-01-1984
		GR 78647 A	27-09-1984
		IE 56902 B	29-01-1992
		IL 69269 A	15-12-1989
		NO 832626 A,B,	23-01-1984
		NZ 204948 A	30-10-1987
		PH 21239 A	21-08-1987
		PL 243134 A	13-08-1984
		PT 77014 A,B	01-08-1983
		US 4673641 A	16-06-1987
		AU 1678783 A	26-01-1984
		JP 2786625 B	13-08-1998
		JP 59118097 A	07-07-1984
		JP 2780961 B	30-07-1998
		JP 8266289 A	15-10-1996
		JP 8294392 A	12-11-1996
WO 9953050 A	21-10-1999	AU 2951499 A	01-11-1999
WO 0044914 A	03-08-2000	NONE	